



*Proceedings of the 4th International Conference and  
Industrial Exhibition on*

# **Dairy Science Park**

**November 1-5, 2017**

**.Abstracts**

**(Incomplete)**

**ISBN-978-969-422-001-7**

## *Editors*

**Rifat Ullah Khan**

**Sania S Qureshi**

**M Subhan Qureshi**

**Mithat Direk**

## *Venue*

**Selçuk Üniversitesi, Konya, Turkey**

## *Co-organizer*

**The University of Agriculture Peshawar-25120, Pakistan**

## *Publisher*

**Engr Irfan Ul Haq Qureshi, President, Dairy Science Park**

**120-Industrial Estate, Peshawar-25000, Pakistan; Mobile/WhatsApp: +92 301 894**

**5994; Email: [i.quraysh@gmail.com](mailto:i.quraysh@gmail.com); Website: <http://dairysciencepark.org.pk/>**

## List of Contents

<b>Item</b>	<b>Page</b>
Foreword	3
Editorial Committee	4
Organizing Committee	5
List of Abstracts	9
Author Index	
Partner Organizations	

## Foreword

Welcome to Konya the city of Mevlana Rumi in Turkey to participate in the Fourth International Conference and Industrial Exhibition on Dairy Science Park scheduled for 1-5 November, 2017, at Selcuk University.

The event is continuation of the series held during November 2011, 2013 and 2015 at Peshawar, Pakistan. The three conferences focused on rehabilitation of the postflood Dairy Sector; Halal meat export potential of the Region and entrepreneurship based hygienic food production respectively. The fourth event of the series is being held with the theme “Achieving Food Security through Entrepreneurship Development and Biorisk Management”.

Pakistan’s Khyber Pakhtunkhwa and FATA are rich in livestock resources valuing US \$10 billion; however, these could neither provide good economic return to producers nor quality food to consumers because of poor management. Similar wastage of natural resources is being observed throughout the world. Dairy Science Park addressed this issue through academic, developmental, business, and marketing interventions. The model has got some notable achievements and further developments are in pipeline. The United Nations have accepted DSP as #SDGAction9671 and it may prove a model for bringing prosperity to the people of the developing nations.

We expect rich participation of people from Academia, Industry, Civil Society, Government and Development Agencies, from various parts of the World. It is hoped that this event will prove as a great opportunity for Academia-Government-Industry linkages and a union between the people of East and West on the historic and spiritual city of Konya.



**Prof M Subhan Qureshi, Co-Chair**  
Email: [drmsqureshi@aup.edu.pk](mailto:drmsqureshi@aup.edu.pk)  
Tel/WhatsApp +92 300 587 7933



**Dr Mithat Direk, Conf Chair**  
Email: [mdirek@selcuk.edu.tr](mailto:mdirek@selcuk.edu.tr)  
Tel/WhatsApp +90 533 577 2651

## Editorial Committee

1. Rifat Ullah Khan, Publication Secretary DSP 2017-Turkey/Assistant Professor (Veterinary Physiology), University of Agriculture, Peshhawar, Pakistan; Email [rukhan@aup.edu.pk](mailto:rukhan@aup.edu.pk)
2. Sania S Qureshi, Editorial Secretary DSP 2017-Turkey/Postgraduate Scholar (Veterinary Pathology) , University of Agriculture, Peshhawar, Pakistan; Email [saniasq@gmail.com](mailto:saniasq@gmail.com)
3. M Subhan Qureshi, Conference Co-Chair/Chief Patron Dairy Science Park, University of Agriculture, Peshhawar, Pakistan; Email [drmsqureshi@aup.edu.pk](mailto:drmsqureshi@aup.edu.pk)
4. Mithat Direk, Conference Chair, Selcuk University, Konya, Turkey, Email: [mdirek@selcuk.edu.tr](mailto:mdirek@selcuk.edu.tr)

## Oganizing Committee

### Honorary Planning Committee

1. Mr Akram Khan Durrani, Minister for Housing and Works, Government of Pakistan, Islamabad
2. Mr Salim Saifullah Khan, Senator, House No.3, Street No. 61,F-6/3, Islamabad
3. Dr. Mukhtar Ahmed, Chairman, Higher Education Commission of Pakistan
4. Mr. Inayatullah Khan, Senior Minister for Local Government, Elections and Rural Development, Government of Khyber Pakhtunkhwa Peshawar, Pakistan
5. Mr Muzaffar Saeed, Minister Finance Department, Government of KP, Peshawar.
6. Mr. Mohibullah Khan, Special Assistant to Chief Minister KP on Livestock and Fisheries, Peshawar, Pakistan
7. Mr. Arif Yousaf, Chairman Task Force on SDGs/Special Assistant to Chief Minister KP on Law and Parliamentary Affairs, Peshawar, Pakistan
8. Prof.Dr. Zahoor Ahmad Swati, Vice Chancellor, The University of Agriculture, Peshawar, Pakistan
9. Prof Dr Ghazala Yasmeen, Vice Chancellor, Women University Mardan
10. Prof.Dr. Mustafa Şahin, President, Selçuk University, Konya, Turkey
11. Prof Peter Wynn, Charles Sturt University, NSW, Australia
12. Ms Nasreen Khattak, Member Provincial Assembly Khyber Pakhtunkhwa, Peshawar.
13. Prof.Dr. Muhammad Sarwar, Vice Chancellor, Gomal University, Dera Ismail Khan, Pakistan
14. Prof.Dr. Mohammad Hafizullah, Vice Chancellor, Khyber Medical University, Peshawar, Pakistan
15. Prof.Dr. Birol Dağ, President, Konya Technopark, Selçuk University, Konya, Turkey
16. Prof.Dr. Tahir Balevi, Vice President, Konya Technopark, Selçuk University, Konya, Turkey
17. Haji Muhammad Afzal, President KP Chamber of Commerce and Industries, Peshawar, Pakistan
18. Prof Khan Bahadar Marwat, Vice Chancellor SBBU, Sheringal, Dir, KP, Pakistan
19. Dr Melissa Finley, Sandia National Laboratories, New Mexico, USA
20. Dr Muhammad Arshad, President Pak Vet Med Council Islamabad/Pak Assoc Adv Sci. Lahore
21. Dr Habib ur Rahman, President Pakistan Veterinary Medical Association KP, Peshawar.
22. Syed Jamal ud Din Shah, Secretary Local Government, GoKP, Peshawar.
23. Dr Khalid Khan, Additional Secretary, HED, Government of KP, Peshawar
24. Mr Shaukat Ali Yousafzai, Additional Secretary, Department of Agriculture, GoKPk, Peshawar.
25. Prof Talat Naseer Pasha, Vice Chancellor UVAS, Lahore, Pakistan
26. Prof Dr Muhammad Tahir Shah, Vice Chancellor, FATA University, Peshawar.
27. Prof Rai Niaz Ahmad, Vice Chancellor PMAS Arid Agricultural University Rawalpindi, Pakistan.
28. Prof Iqrar Ahmad Khan, Vice Chancellor, University of Agriculture, Faisalabad, Pakistan

29. Prof Khuda Bakhsh Mirbahar, Vice Chancellor, SBB-UVAS, Sakrand, Sindh Pakistan

### **Organizing Committee**

1. Prof M Azam Kakar, BUITMS/World Bank, Quetta
2. Malik Niaz Ahmad, Member KP Chamber of Commerce and Industries, Peshawar, Pakistan.
3. Mr. Abdul Rahman Ilyas, Advisor Agribusiness, Economic Development Board, Government of Andhra Pradesh, Hyderabad, India
4. Muhammad Zubair Mughal, CEO, Halal Research Council, 117, G – III, Johar Town Lahore, Pakistan
5. Dr. Muhammad Latif, Adviser (R&D) HEC, Islamabad
6. Prof Dr Hamayun Khan, President AUTA, University of Agriculture, Peshawar
7. Mr. Ahmad Naveed, Director General, Livestock and Dairy Development (Research Wing), Peshawar, Pakistan
8. Mr. Ahmad Waleed Joyan, Sandia National Laboratories, Albuquerque, New Mexico, USA
9. Prof.Dr. Gang Li, Head, DCPAID, Institute of Animal Sciences, CAAS, Beijing, China
10. Prof Muhamad Jamal Khan, Dean FCPS, UA Peshawar
11. Dr Syed Naeem Badshah, Chairman Islamic Studies, Univ Agric, Peshawar
12. Prof Saifullah, Dean Faculty of Crop Protection Sciences, University of Agriculture, Peshawar.
13. Prof Masood Rabbani, Dean FVS, University of Veterinary and Animal Sciences, Lahore
14. Dr Hina Jabeen, Head Microbiology, Women University Mardan
15. Dr Rajwali Khan, PhD Scholar, North-West Agriculture and Forestry University, Yangling, Shaanxi, China
16. Prof Hameedullah Shah, Dean Faculty of Nutrition Sciences, University of Agriculture, Peshawar.
17. Prof Miraj ul Islam Zia, Dean Faculty of Islamic and Oriental Studies, University of Peshawar.
18. Prof Noor Pao Khan, Dean Faculty of Rural and Social Sciences, University of Agriculture, Peshawar.
19. Mr Naimatullah Khan, Director Finance, University of Agriculture, Peshawar.
20. Prof M Saleem Chaudhry, Pak Assoc Adv Science, Lahore.
21. Mr. Sahibzada Shafiullah, School of Animal & Veterinary Sciences, Charles Sturt University, Wagga Wagga, Australia
22. Mr. Sher Muhammad, Director General, Livestock and Dairy Development (Extension Wing), Peshawar, Pakistan
23. Mr Israr Abbas, Pakistan Science Foundation, Islamabad.
24. Dr Aftab Ahmad, President National Academy of Young Scientists, Islamabad, Pakistan.
25. Mr Javed Khattak, Provincial Chief, Small and Medium Development Authority, Peshawar, Pakistan.
26. Prof Shaukat Hussain, Chairman Plant Pathology, University of Agriculture, Peshawar
27. Prof Imtiaz Ali Khan, Entomology Department, University of Agriculture, Peshawar

28. Mr. Tawheed Ali Azimee, Faculty of Agriculture, Nangarhar University, Jalalabad, Afghanistan
29. Dr. Zia Ul Haq, Director, ORIC, Khyber Medical University, Peshawar, Pakistan
30. Dr Muhammad Inam, HoD Animal Sciences, SBBU, Sheringal, Dir Upper, Pakistan
31. Prof Ahrar Khan, Dean FVS, University of Agriculture, Faisalabad, Pakistan
32. Mr Rizwan Ahmad, Director P&D, University of Agriculture, Peshawar.
33. Prof Umer Sadique, Chairman Animal Health, The University of Agriculture, Peshawar, Pakistan
34. Ms Nighat Amjad, AIFA Environmental Services, Dubai, UAE
35. Dr Irfan Khattak, Abdul Wali Khan University Mardan

### Scientific Committee

1. Prof.Dr Zafar Iqbal, Faculty of Veterinary Sciences, University of Agriculture, Faisalabad
2. Prof Anas Sarwar, FVS, University of Agriculture, Faisalabad, Pakistan.
3. Dr. Muhammad Ovais Omer, Chairman Pharmacology, University of Veterinary and Animal Sciences, Lahore
4. Dr Murad Ali Khan, Secretary AUTA, University of Agriculture, Peshawar.
5. Dr Zubia Masood, Jinnah University for Women, Karachi, Pakistan
6. Prof.Dr. Ahmet Güner, Dean, Faculty of Veterinary, University of Selçuk, Konya, Turkey
7. Prof.Dr. Mustafa Karakaya, Faculty of Agriculture, Selçuk University, Konya, Turkey
8. Prof.Dr. Nihat Akın, Head, Department of Food Science, Faculty of Agriculture, Selçuk University, Konya, Turkey
9. Prof.Dr. Cevat Aydın, Dean, Faculty of Agriculture, Selçuk University, Konya, Turkey
10. Dr Saim Boztepe, Dept Anim Sci, Agric Faculty, Selcuk University, Konya
11. Prof Ali Akmaz, Faculty of Veterinary, University of Selcuk, Konya, Turkey
12. Prof Cennet Oguz, Agric Economics, Selcuk University, Konya, Turkey
13. Dr Komariah, Agroclimatology, Sebelas Maret, University of Indonesia
14. Haroun Chnchouni, Ecology, University of Tebessa, Algeria
15. Prof Yusuf Konca, Zootečni Dept, Univ Erciyes, Kayseri, Turkey
16. Prof Zafer Ulutas, Dept Anim Prod, Omer Halis Demir Univ, Nigde, Turkey
17. Prof Darlina Md Naim, Biol Sci, Univ Sains Malaysia, Penang, Malaysia
18. Prof Mehmet Koyunu, Zootečni Dept, University of Uludag, Bursa, Turkey
19. Prof Feyzi Ugur, Zootečni Dept, Univ Canakkale, Turkey
20. Prof Shazia Shafique, Plant Pathology, University of the Punjab, Lahore
21. Prof Sobia Shafique, Mycology and Plant Pathology, Univ Punjab, Lahore
22. Prof Felix Arion, Univ Agric Sci & Veterinary Medicine, Romania
23. Dr Osman Ozbek, Department of Agric Mech & Techn, Selcuk Univ, Konya
24. Prof AM Ahmad, Agribusiness & Applied Economics, Tanta Univ Egypt
25. Prof Prabang Setyono, Env Sci, Sebelas Maret University, Indonesia
26. Prof Widyatmani Sih Dewi, Soil Sci, Sebelas Maret University, Indonesia
27. Dr Halils Simsek, Agric & Biosystems Engg, Noth Dekota State Univ, USA
28. Dr Syed Muhammad Suhail, PRO, Dairy Science Park, Peshawar
29. Prof Arfan Yousaf, Dean FAHVS, PMAS Arid Agricultural University, Rawalpindi, Pakistan
30. Dr Tariq Ali Khattak, Animal Health Department, UA Peshawar

31. Dr Farhan Anwar Khan, Animal Health Department, UA Peshawar
32. Dr Nazir Ahmad Khan, AN Department, UA Peshawar
33. Dr Niaz Ali, IBMS, Khyber Medical University, Peshawar
34. Dr Khalid Khan, PhD Scholar ABG, UA Peshawar
35. Dr Hamayun Khan, Assistant Professor Anatomy, UA Peshawar
36. Dr Ikramullah Khan, Livestock and Dairy Development Department, Bannu
37. Dr Shakirullah, Assistant Professor GCVS, Gomal University DI Khan
38. Dr Muhammad Sohail Sajid, Associate Professor Parasitology, University of Agriculture Faisalabad

### **Management Committee**

1. Dr Mithat Direk, Chair DSP 2017, Konya, Email: [mdirek@selcuk.edu.tr](mailto:mdirek@selcuk.edu.tr) , Tel: +90 533 577 2651
2. Prof M Subhan Qureshi, Co-Chair DSP 2017, Konya, Email: [drmsqureshi@aup.edu.pk](mailto:drmsqureshi@aup.edu.pk); Tel +92 300 587 7933
3. Conference Secretaries
4. General: Irfan ul Haq Qureshi, DSP, Tel +92 301 894 5994, Email [i.quraysh@gmail.com](mailto:i.quraysh@gmail.com)
5. Development: Prof Sarzamin Khan, [dr.zaminaup@gmail.com](mailto:dr.zaminaup@gmail.com)
6. Publication: Dr Rifat Ullah Khan [rifatullahkhan@gmail.com](mailto:rifatullahkhan@gmail.com)
7. Scientific: Dr Shakoor Ahmad, UAP, Tel 0345 345 5456, Email [shakoor.ahmad@aup.edu.pk](mailto:shakoor.ahmad@aup.edu.pk)
8. Editorial: Ms Sania Qureshi; [saniasq@gmail.com](mailto:saniasq@gmail.com)
9. Commercial: Rashid Aman, SMEDA; +92 333 911 7278; Email: [mrashid@smeda.org.pk](mailto:mrashid@smeda.org.pk);
10. Academic: Nasir Shah HEC; Tel: +92 300 596 0299; Email: [nrshah@hec.gov.pk](mailto:nrshah@hec.gov.pk)
11. Scientific Coordinator (Turkey): Aamir Iqbal Khattak; WhatsApp +92 344 957 2556



## List of abstracts

(Categories: AH=Animal Health; AP=Animal Production; FSE=Food Security and Ethics; MT=Meat Technology; DT=Dairy Technology; BRM=Biorisk Management; PH=Public Health; QC=Quality Control; ED=Entrepreneurship Development)

Code	Category	Title/authors/corresponding email	Institution	Presentation
001	BRM	SEROPREVALENCE AND POTENTIAL RISK FACTORS ASSOCIATED WITH BOVINE BRUCELLOSIS UNDER DIVERSE PRODUCTION SYSTEMS IN PAKISTAN Aman Ullah Khan*, Iahtasham Khan, Muhammad Younus, Syed Ehtisham-Ul-Haque, Usman Waheed, Falk Melzer, Hosny El-Adawy, Heinrich Neubauer, Muhammad Asif, Idrees, <a href="mailto:amanullah.khan@uvas.edu.pk">*amanullah.khan@uvas.edu.pk</a>	UVAS Lahore, Pakistan	Oral
002	BRM	CRIMEAN CONGO HEMORRHAGIC FEVER (CCHF) IN THE SCENARIO OF DSP-BIORISK MANAGEMENT IN PAKISTAN P.R. Nasrullah, M.A. Kakar*, Azizullah, M.S. Qureshi and Mithat Direk <a href="mailto:adaqul@yahoo.com">adaqul@yahoo.com</a>	AUP Peshawar	
003	BRM	QUALITY ASSESMENT OF PASTEURIZED MILK THROUGH THE INCIDENCE OF SPORE FORMING BACTERIA IN PASTEURIZED MILK Muhammad Asim Ikram*, Imran Javed, Saima Inayat, Khalid Javed, M. Junaid, H. Arslan Amin, Usman Mir Khan <a href="mailto:asim.ikram@uvas.edu.pk">*asim.ikram@uvas.edu.pk</a>	UVAS Lahore	
004	ED	CHALLENGES AND FUTURE SCOPE OF OSTRICH (STRUTHIO CAMELIUS) FARMING IN PAKISTAN Muhammad Fiaz Qamar*, Muhammad Arfan Zaman, and Kazim Ali <a href="mailto:fiaz.qamar@uvas.edu.pk">*fiaz.qamar@uvas.edu.pk</a>	UVAS Lahore	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
005	BRM	ESTIMATION OF COMPOSITION, ADULTERATION AND MICROBIAL LOAD IN RAW MILK SAMPLES COLLECTED FROM DAIRY FARMS AND URBAN AREAS OF LAHORE DISTRICT, PAKISTAN Atta Muhammad Arif, Muhammad Ali* <a href="mailto:Muhammadali.uvas@gmail.com">*Muhammadali.uvas@gmail.com</a>	UVAS Lahore	Oral
006	AH	ANTHELMINTIC AND ANTIPROTOZOAL ACTIVITIES OF <i>CALOTROPIS PROCERA</i> UNDER FIELD CONDITIONS Muhammad. Ashraf*, A. Javeed, M. Adil Rasheed, Qamar Niaz and Mukhtar Ahmad <a href="mailto:drashraf2001@uvas.edu.pk">*drashraf2001@uvas.edu.pk</a>	UVAS Lahore	Oral
007	BRM	TOXICOLOGICAL EFFECTS OF AFLATOXIN B1 ON GROWTH PERFORMANCE, HUMORAL IMMUNE RESPONSE AND BLOOD PROFILE OF JAPANESE QUAIL Muhammad Younus*, Sakhra Mahmood, Asim Aslam, Aftab Ahmed Anjum, Aman Ullah Khan, Asif Idrees, M.L. Sohail <a href="mailto:younusrana@hotmail.com">*younusrana@hotmail.com</a>	UVAS Lahore	Oral
008	AH	DISPOSITION KINETICS OF CIPROFLOXACIN AFTER INTRAMUSCULAR ADMINISTRATION IN <i>LOHI</i> SHEEP Zahid Iqbal*, Ijaz Javed, Riaz Hussain, Mudassar Mohiuddin <a href="mailto:zahid1.iqbal1@gmail.com">*zahid1.iqbal1@gmail.com</a>	Isra University	Oral
009	AP	EFFECTS OF DIETARY VITAMIN C ON ARSENIC INDUCED TOXICITY ON SEMEN QUALITY, REPRODUCTIVE HORMONES AND HISTOPATHOLOGY OF TESTES IN TEDDY GOAT BUCKS Muhammad Zubair*, Maqbool Ahmad, Mushtaq Ahmad, Saqib Umar, Shujait Ali <a href="mailto:drzubairabbasi@gmail.com">*drzubairabbasi@gmail.com</a>	University of Ponch, Azad Kashmir	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
010	MT	CHARACTERISTICS OF MEAT EMULSION SYSTEMS WITH BEEF REPLACED BY SOY PRODUCTS Neslihan Polat, Mustafa Karakaya, Ali Samet Babaoğlu*, Kübra Ünal * <a href="mailto:asmtbb@gmail.com">asmtbb@gmail.com</a>	Selcuk University, Konya, Turkey	Oral
011	QC	A REVIEW ON AFLATOXIN M1 PREVALENCE IN MILK AND RECENT UPDATES ON DECONTAMINATION STRATEGIES Amir Ismail*, Muhammad Riaz, Saeed Akhtar * <a href="mailto:amirismail@bzu.edu.pk">amirismail@bzu.edu.pk</a>	Bahauddin Zakariya University, Multan, Pakistan	Oral
012	DT	COLOSTRUM WHEY: A POTENTIAL INGREDIENT FOR DAIRY SECTOR IN NUTRITIONAL ASPECTS Maryam Batool*, Saima Inayat, Muhammad Asim Ikram * <a href="mailto:maryambatool17@gmail.com">maryambatool17@gmail.com</a>	UVAS, Lahore	Oral
013	BRM	DIFFERENTIAL GROWTH OF SALMONELLA PULLORUM AND SALMONELLA GALLINARUM USING MODIFIED SS AGAR AND RV MEDIUM Bakhtawar Maqbool*, Sajjad-ur-Rahman * <a href="mailto:bakhtawardogar@yahoo.com">bakhtawardogar@yahoo.com</a>	University of Agriculture, Faisalabad, Pakistan	Oral
014	AH	COMPARATIVE RESILIENCE OF TEDDY AND BEETAL GOAT BREEDS AGAINST ARTIFICIAL INFECTION OF HAEMONCHUS CONTORTUS THROUGH IMMUNOGLOBULINS AND HISTAMINE Asim Shamim*, Muhammad Sohail Sajid, Muhammad Nisar Khan, Muhammad Saqib, Muhammad Arfan Zaman, Muhammad Imran, Hafiz Muhammad Rizwan * <a href="mailto:asimshamimuajk@gmail.com">asimshamimuajk@gmail.com</a>	University of Poonch Rawalakot Azad Kashmir, Pakistan	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
015	AH	HIGH PROTEIN DIET INDUCES B-CELL REGENERATION AND RESTORES NORMAL GLYCAEMIA IN ALLOXAN INDUCED DIABETIC RATS Humaira Muzaffar*, Muhammad Naeem Faisal, Tanweer Khaliq, Junaid Ali Khan, Bilal Aslam, Aisha Mahmood <a href="mailto:2007ag1556@uaf.edu.pk">*2007ag1556@uaf.edu.pk</a>	University of Agriculture Faisalabad, Pakistan	Poster
016	AH	ANTIMICROBIAL POTENTIAL OF BOVINE LACTOFERRIN AGAINST FOODBORNE PATHOGENS Ishtiaque Ahmad*, Ammarah Khatoon, Muhammad Ayaz, Muhammad Asim Ikram, Muhammad Junaid, Usman Mir Khan, Nisar Ahmad <a href="mailto:ishtiaque@uvas.edu.pk">*ishtiaque@uvas.edu.pk</a>	UVAS Lahore, Pakistan	Oral
017	MT	THE EFFECTS OF KEFIR IN TENDERIZING CHICKEN AND TURKEY BREAST MEAT Kübra Ünal*, Ali Samet Babaoğlu, Mustafa Karakaya <a href="mailto:ulusoy_kubra@hotmail.com">*ulusoy_kubra@hotmail.com</a>	Selçuk University, Konya, Turkey	Poster
018	AH	COMPARATIVE EFFICACY OF SYNCHRONIZATION PROTOCOLS (PROGESTERONE BASED OR FREE) ON CONCEPTION RATE IN CAPRINE SPECIES Maqbool Ahmad*, Umair Asharaf, Shafia Tehseen Gul, Rehana Kousar, Ijaz Ahmad, Khurram Ashfaq <a href="mailto:drmaqbool_ahmad@yahoo.com">*drmaqbool_ahmad@yahoo.com</a>	University of Agriculture, Faisalabad, Pakistan	Oral
019	ED	WATER PRODUCTIVITY UNDER GOVERNMENT AND FARMERS CONTROLLED CANAL IRRIGATION SYSTEMS Muhammad Jamal Khan*, Rabnawaz and Amanullah <a href="mailto:mjkhan@aup.edu.pk">*mjkhan@aup.edu.pk</a>	The University of Agriculture Peshawar, Pakistan	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
020	BRM	SYNTHETIC POLYMORPHIC PEPTIDES: A STEP FORWARD FOR DIAGNOSIS OF TYPE-SPECIFIC TOXOPLASMOSIS IN HUMANS AND ANIMALS H.M. RIZWAN*, D. ARRANZ-SOLÍS, MS SAJID, J.P. SAEIJ * <a href="mailto:dr.hmrizwan@yahoo.com">dr.hmrizwan@yahoo.com</a>	The University of Agriculture Faisalabad, Pakistan	Oral
021	AP	FASCIOSIS: A MAJOR THREAT TO FOOD SECURITY AND ECONOMIC LOSSES ALL AROUND THE WORLD Hafiz Muhammad Rizwan*, Muhammad Sohail Sajid, Muhammad Saqib, Zafar Iqbal * <a href="mailto:dr.hmrizwan@yahoo.com">dr.hmrizwan@yahoo.com</a>	The University of Agriculture Faisalabad, Pakistan	Poster
022	AP	EFFECT OF BREED ON CARCASS TRAITS OF KUNDHI AND NILI RAVI BUFFALALOES Rameez Raja Kaleri*, Hubdar Ali Kaleri, Muhammad Siddique Zardari * <a href="mailto:rameezkaleri@gmail.com">rameezkaleri@gmail.com</a>	Sindh Agriculture University, Tandojam, Pakistan	Oral
023	ED	DAIRY SCIENCE PARK–EMPLOYMENT OPPORTUNITIES FOR THE YOUTH Sania Subhan Qureshi*, Irfan ul Haq Qureshi, Sarzamin Khan and M Subhan Qureshi * <a href="mailto:saniasq@aup.edu.pk">saniasq@aup.edu.pk</a>	The University of Agriculture, Peshawar, Pakistan	Poster
024	DT	EFFECT OF RIPENING CONDITIONS ON PHYSICO-CHEMICAL PROPERTIES AND SURVIVAL OF PROBIOTIC MICROORGANISMS IN PROBIOTIC CHEDDAR CHEESE Usman Mir Khan*, Fizzah Ashfaq, Arslan Amin Bhatti, Ishtiaque Ahmad, Saima Inayat, Muhammad Ayaz, Nisar Ahmad * <a href="mailto:usmanmirkhan@yahoo.com">usmanmirkhan@yahoo.com</a>	UVAS, Lahore, Pakistan	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
025	DT	PHYSICO-CHEMICAL AND NUTRITIONAL PROPERTIES OF CHEDDAR CHEESE MADE FROM CITRUS RETICULATA BLANCO CRUDE FLOWERS EXTRACT (CFE) Usman Mir Khan*, Ishtiaque Ahmad, Saima Inayat, Muhammad Ayaz, Nisar Ahmad, Arslan Amin Bhatti * <a href="mailto:usmanmirkhan@yahoo.com">usmanmirkhan@yahoo.com</a>	UVAS Lahore, Pakistan	
026	DT	ENHANCEMENT OF OXIDATIVE STABILITY OF IRON ENRICHED BUTTER OIL WITH NATURAL ANTIOXIDANTS Ikam Ullah, Ishtiaque Ahmad, Usman Mir Khan*, Nisar Ahmad * <a href="mailto:usmanmirkhan@yahoo.com">usmanmirkhan@yahoo.com</a>	UVAS Lahore, Pakistan	
027	ED	STRENGTHENING COMMUNITY RADIO FOR EFFECTIVE CLIMATE CHANGE ADAPTATION STRATEGIES FOR CLIMATE SMART AGRICULTURAL PRODUCTION IN AFRICA Wilson Okaka* and Jennifer Apil <a href="mailto:nupap2000@yahoo.com">nupap2000@yahoo.com</a>	Makerere University, Uganda	Poster
028	ED	PUBLIC AWARENESS COMMUNICATION CAMPAIGNS FOR DIFFUSION OF FOOD AND NUTRITION SECURITY POLICY INFORMATION SERVICES FOR SUSTAINABLE LIVELIHOODS IN UGANDA Wilson Okaka* and Jennifer Apil * <a href="mailto:nupap2000@yahoo.com">nupap2000@yahoo.com</a>	Kyambogo University, Uganda	Oral
029	ED	AN INVESTIGATION ON CHEMICAL COMPOSITION AND NUTRITIONAL EVALUATIONS OF SOME NATURALLY PRODUCE FISHERIES RESOURCES OF PAKISTAN Zubia Masood*, Nelofer Jamil, Sarzamin Khan * <a href="mailto:zubiamasood12@gmail.com">zubiamasood12@gmail.com</a>	Jinnah University for Women, Karachi, Pakistan	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
030	AD	<p>EPIDEMIOLOGY AND CONTROL OF GASTROINTESTINAL FAUNA OF EUROPEAN ALBINO RABBITS OF FAISALABAD DISTRICT, PUNJAB, PAKISTAN</p> <p>Muhammad Sohail Sajid*, Waqas Altaf, Muhammad Kasib Khan Muhammad Saqib  <a href="mailto:drsohailuaf@hotmail.com">*drsohailuaf@hotmail.com</a></p>	UAF Pakistan	Oral
031	BRM	<p>COMPARATIVE EVALUATION OF MICROBIOLOGICAL MILK SAFETY STANDARDS EXISTING IN FAISALABAD</p> <p>Nabila Yaqoob, Ahsan Naveed*, Sajjad Ur Rahman, Syeda Wajiha Hassan, M. Farooque Hassan  <a href="mailto:ahsannaveed.uaf@gmail.com">*ahsannaveed.uaf@gmail.com</a></p>	UAF Pakistan	Poster
032	AH	<p>HIGH OMEGA-6/OMEGA-3 RATIO IN MARGARINE INDUCES INTESTINAL INFLAMMATION AND OXIDATIVE STRESS RESPONSE IN WISTER RAT</p> <p>Aisha Mahmood*, Muhammad Naeem Faisal, Tanweer Khaliq, Junaid Ali Khan, Faqir Muhammad, Humaira Muzaffar, Farooq Azam and Maha Saleem  <a href="mailto:2007ag1551@uaf.edu.pk">*2007ag1551@uaf.edu.pk</a></p>	UAF Pakistan	Oral
033	ED	<p>EFFECT OF GAMMA-IRRADIATION ON THE QUALITY CHARACTERISTICS OF BROWN RICE BASED WEANING FOOD</p> <p>Awsi Jan*, Dr. Monika Sood, P.R.Hussain, Lubna Masoodi, Ambreena, Nadia Bashir  <a href="mailto:awsijan@gmail.com">*awsijan@gmail.com</a></p>		
034	BRM	<p>PREVALENCE OF <i>ESCHERICHIA COLI</i> O157: H7 IN BEEF MEAT IN WESTERN ALGERIA</p> <p>Barka M Salih* Benyoub Noreddine  <a href="mailto:bm_salih@hotmail.com">bm_salih@hotmail.com</a></p>	Route de la Rocade, Algeria	Poster
035	BRM	<p>Risk characterization of antibiotic resistance of Salmonella isolated from raw meat in Penang, Malaysia; Hafiz NidaUllah*, Mohd Omar Abdul Kadir, Rosma Ahmad, Widad Fadhullah &amp; Saima Sohni; <a href="mailto:hafiz.nidaullah@daad-alumni.de">hafiz.nidaullah@daad-alumni.de</a></p>	Universiti Sains Malaysia	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
036	AP	COMPARATIVE EVALUATION OF NIGELLA SATIVA AND LINCOMYCIN AS GROWTH PROMOTERS IN COMMERCIAL BROILER Munnaza Shareef, Muhammad Naeem Faisal, Aisha Mahmood, Humaira Muzaffar, JazibHussain*, Samia Ali, Muhammad Jamal, Farooq Azam <a href="mailto:jazibhussain04@gmail.com">jazibhussain04@gmail.com</a>	UAF Pakistan	Oral
037	DT	ESTIMATION OF MILK PROTEASE ACTIVITY FROM UNINFECTED MAMMARY GLANDS OF NILI-RAVI BUFFALOS , SAHIWAL AND CROSS-BREED COWS OF PAKISTAN Razia Kausar*, Amjad Hameed, and Zafar Iqbal Qureshi <a href="mailto:razia_uaf@hotmail.com">*razia_uaf@hotmail.com</a>	UAF	Oral
038	ED	APPLICATIONS OF PULSE ELECTRIC FIELD IN DAIRY TECHNOLOGY Saleem Malik*, Lubna Masoodi , Awsi Jan <a href="mailto:malik.saleem72@gmail.com">*malik.saleem72@gmail.com</a>	Sher-e-Kashmir University, Pakistan	Poster
039	AP	EFFECT OF MENTHA PIPERITA (PEPPERMINT) EXTRACT AND ITS JUICE ON EGG QUALITY TRAITS DURING DIFFERENT STORAGE TIME IN LAYING HENS Abdur Rahman, E. Eren Gültepe, Cangir Uyarlar, I. Sadi Çetingül, Aamir Iqbal* , Ismail Bayram <a href="mailto:aamir_vet@yahoo.com">aamir_vet@yahoo.com</a>	UVAS, Lahore-Pakistan	Oral
040	AP	THERMAL AND NUTRITIONAL STRESS IN EWES AND DOES: EFFECTS ON EMBRYO DEVELOPMENT AND EMBRYONIC LOSSES Dehouegnon Jerry Agossou*, Uğur Serbester, Nazan Koluman <a href="mailto:bermondjerry@gmail.com">bermondjerry@gmail.com</a>	Çukurova University	Oral



Code	Category	Title/authors/corresponding email	Institution	Presentation
041	AP	COMPARISON OF LINCOMYCIN AND TETRACYCLINE AS ANTIBIOTIC GROWTH PROMOTER IN COMMERCIAL BROILER Muhammad Jamal, Muhammad Naeem Faisal, Hamza Nazeer*, Humaira Muzaffar, Aisha Mahmood, Jazib Hussain, Jawad Aslam, Hira Javed <a href="mailto:hamzanazir56@yahoo.com">hamzanazir56@yahoo.com</a>	UAF, Faisalabad, Pakistan	Poster
042	PH	PHYTOREMEDIATION: AN INSPIRING AND ECONOMICAL FOR THE TREATMENT OF IRRIGATION WATER Hina Jabeen*, Aqsa Gilani, Fatima Waleed, Mushtaq Ahmad, Rakhshinda Sadiq, Akhtar Rasool <a href="mailto:hinajabeen2@gmail.com">hinajabeen2@gmail.com</a>	Women University Mardan, KPK Pakistan	Oral
043	ED	PRODUCTION AND IMMOBILIZATION OF ENDOGLUCANASES FROM MUTANT DERIVATIVE OF SUPER KOJI (ASPERGILLUS ORYZAE) ON NANOPOROUS SUPPORT Neelam Zeb <sup>1,2</sup> , Dr. Muhammad Hamid Rashid <sup>2</sup> <a href="mailto:neelamzeb@ymail.com">neelamzeb@ymail.com</a>	Women University Mardan	Oral
044	ED	EFFECT OF ANTIOXIDANT, PACKAGING MATERIAL AND STORAGE PERIODS ON THE CHEMICAL COMPOSITION OF ALMOND KERNEL Lubna masoodi*, Dr A.H Rather, Awsi Jan <a href="mailto:lubna.masudi@gmail.com">lubna.masudi@gmail.com</a>	Shere-e- Kashmir University of Agricultural Sciences and Technology Srinagar Jammu and Kashmir	Oral
045	PH	AN ANALYSIS OF CORRELATION BETWEEN POOR FOOD EDUCATION AND DENTAL CARIES IN PESHAWAR Maria Nizam* <a href="mailto:Marianizam65@gmail.com">Marianizam65@gmail.com</a>	Gandhara University Peshawar, Pakistan	Oral

Code	Category	Title/authors/corresponding email	Institution	Presentation
046	AP	MONOGLYCERIDES AN ALTERNATIVE TO ANTIBIOTIC GROWTH PROMOTERS IN COMMERCIAL BROILER Samia Ali, Muhammad Naeem Faisal, Aisha Mahmood, Humaira Muzaffar, Muhammad Jawad Aslam* <a href="mailto:jawadaslamch@gmail.com">jawadaslamch@gmail.com</a>	UAF, Pakistan	Oral
047	AP	EFFECT OF FENUGREEK SEED SUPPLEMENTATION ON EGG QUALITY AND PRODUCTION PERFORMANCE OF SPENT LAYERS N. Chand.*, F.Wahab, M.S. Qureshi and R.U. Khan <a href="mailto:drleeze@yahoo.com">drleeze@yahoo.com</a>	AUP	Oral
048	ED	DETERMINATION OF RED AND WHITE MEAT CONSUMPTIONS OF ADULT INDIVIDUALS LIVING IN AKŞEHİR Nazik Meziyet DİLEK*, Ezgi TOPTAŞ BIYIKLI, Ali Emrah BIYIKLI <a href="mailto:meziyettemel@hotmail.com">meziyettemel@hotmail.com</a>	Selcuk University, Turkey	Poster
049	ED	IDENTIFICATION OF THE PREFERENCES OF HEALTHY ADULTS ABOUT RED AND WHITE MEAT CONSUMPTION Nazik Meziyet DİLEK*, Ali Emrah BIYIKLI, Ezgi TOPTAŞ BIYIKLI <a href="mailto:meziyettemel@hotmail.com">meziyettemel@hotmail.com</a>	Selcuk University, Turkey	Poster
050	PH	CONJUGATED LINOLEIC ACID IN ANIMAL SOURCE FOODS AND ITS EFFECT ON HEALTH Nazik Meziyet DİLEK*, Mustafa KARAKAYA <a href="mailto:meziyettemel@hotmail.com">meziyettemel@hotmail.com</a>	Selcuk University, Turkey	Poster
051	PH	IDENTIFICATION OF PROCESSED MEAT PRODUCTS CONSUMPTION OF HEALTHY ADULT INDIVIDUALS LIVING IN AKŞEHİR Nazik Meziyet DİLEK*, Ezgi TOPTAŞ BIYIKLI, Ali Emrah BIYIKLI <a href="mailto:meziyettemel@hotmail.com">meziyettemel@hotmail.com</a>	Selcuk University, Turkey	Poster

Code	Category	Title/authors/corresponding email	Institution	Presentation
052	PH	POTENTIAL OF RED MEAT PRODUCTION IN TURKEY Nazik Meziyet DİLEK*, Mustafa KARAKAYA <a href="mailto:meziyettemel@hotmail.com">meziyettemel@hotmail.com</a>	Selcuk University, Turkey	Poster
053	ED	NANOTECHNOLOGY IN AGRICULTURE AND FOOD INDUSTRY: AN OVERVIEW OF APPLICATIONS AND CHALLENGES Rakhshinda Sadiq* and Hina Jabeen <a href="mailto:rakhshinda_sadiq@yahoo.com">rakhshinda_sadiq@yahoo.com</a>	Women University Mardan, Mardan	Oral
054	ED	THREATENED WILD MORELS AND MOUNTAIN LIVELIHOODS IN NORTH WEST PAKISTAN Saadia Hanif* <a href="mailto:saadia.hanif@live.com">saadia.hanif@live.com</a>	Consultant-Farm Dynamics Pakistan (pvt) Ltd.	Oral
055	ED	POLITICAL/IDEOLOGICAL STUDY AND ACADEMIC STUDY: SCIENCE,HISTORY AND TEACHING. Sarmad Hussain Tunio* <a href="mailto:Sarmad.hussain@nu.edu.pk">Sarmad.hussain@nu.edu.pk</a>	NU-FAST Karachi	Poster
056	AH	THERAPEUTIC EFFECT OF HERBAL PLANTS EXTRACT IN BROILER CHICKEN CHALLENGED WITH COCCIDIA Subhan Ullah and Umar Sadique* <a href="mailto:druskhhatak@yahoo.com">druskhhatak@yahoo.com</a>	AUP	Oral
057	AH	GENOTOXIC AND TOXICOPATHOLOGICAL EFFECT OF AFLATOXIN B1 IN GRASS CARP (Ctenopharyngodon idella) Hayatullah Khan and Umer Saddique* <a href="mailto:druskhhattak@yahoo.com">druskhhattak@yahoo.com</a>	AUP	Oral
058	AH	ANAEMIA – STILL A MAJOR HEALTH PROBLEM Binish Sarwar Khan* <a href="mailto:Binishasad786@gmail.com">Binishasad786@gmail.com</a>	UAF	Poster

Code	Category	Title/authors/corresponding email	Institution	Presentation
059	BRM	OCCURRENCE OF MYCOBACTERIUM BOVIS IN BOVINE MILK AND CONSUMERS KNOWLEDGE, ATTITUDE AND PRACTICES TOWARD TUBERCULOSIS IN PESHAWAR, PAKISTAN Irfan Khattak* <a href="mailto:irfankhattak@awkum.edu.pk">irfankhattak@awkum.edu.pk</a>	Abdul Wali Khan University, Mardan, Pakistan	Oral
060	AH	TOXINOTYPING OF CLOSTRIDIUM PERFRINGENS INFECTED LAMBS AND KIDS IN DIFFERENT TOPOGRAPHIC AREAS OF KHYBER PAKHTUNKHWA, PAKISTAN Mumtaz Ali Khan* <a href="mailto:Dr_mumtazkhan@yahoo.com">Dr_mumtazkhan@yahoo.com</a>	UVAS Lahore	Oral
061	AH	HEMATO-BIOCHEMICAL ALTERATIONS IN CLOSTRIDIUM PERFRINGENS INFECTED GOATS Mumtaz Ali Khan* <a href="mailto:dr_mumtazkhan@yahoo.com">dr_mumtazkhan@yahoo.com</a>	UVAS, Lahore, Pakistan	
062	AH	QUANTIFICATION, ISOLATION AND PATHOGENOMICS F PATHOGENIC SHIGA TOXIN ESCHERICHIA COLI O157:H7 ALONG THE PRODUCTION AND SUPPLY CHAIN AROUND HUBEI OF CHINA Sher Bahadar Khan* <a href="mailto:sbvetdr@yahoo.com">sbvetdr@yahoo.com</a>	AUP	Oral
063	BRM	ANTIMICROBIAL RESISTANCE OF BACTERIA ASSOCIATED WITH RAW MILK CONTAMINATED WITH ANTIBIOTICS RESIDUES Limya .M. Warsma* <a href="mailto:areejwarsmaq3@gmail.com">areejwarsmaq3@gmail.com</a>	University of Khartoum, Sudan	Oral
064	ED	PROCESSING AND UTILIZATION OF TRADITIONAL DAIRY PRODUCTS IN RAWALAKOT AZAD JAMMU AND KASHMIR, PAKISTAN Imtiaz Hussain* <a href="mailto:imtiazupr@yahoo.com">imtiazupr@yahoo.com</a>	University of Poonch, Azad Kashmir	Poster

Code	Category	Title/authors/corresponding email	Institution	Presentation
065	ED	VALUE ADDITION AND PROCESSED PRODUCTS OF UNDERUTILIZED FRUITS FOR FOOD SECURITY IN AZAD JAMMU AND KASHMIR, PAKISTAN Imtiaz Hussain* <a href="mailto:imtiazuپر@yahoo.com">imtiazuپر@yahoo.com</a>	University of Poonch, Azad Kashmir	Poster
066	AH	HYGIENE AND DAIRY MANAGEMENT PRACTICE DURING MILKING AT RAWALAKOT AZAD JAMMU AND KASHMIR, PAKISTAN Imtiaz Hussain* <a href="mailto:imtiazuپر@yahoo.com">imtiazuپر@yahoo.com</a>	University of Poonch, Azad Kashmir	Oral
067	AH	NUTRACEUTICAL APPLICATIONS OF ZINC, MANGANESE AND COPPER IN THE METABOLISM OF THE RUMEN AND IMMUNE FUNCTION Habib-Ur-Rehman* <a href="mailto:hur_2085qta@yahoo.com">hur_2085qta@yahoo.com</a>	University of Balochistan (UoB), Quetta, Pakistan	Oral
068	ED	REUSE OF WASTEWATER IN AGRICULTURE RECOVERED FROM WASTEWATER TREATMENT SYSTEMS; Khush Bakht Andleeb* <a href="mailto:kbandleeb@gmail.com">kbandleeb@gmail.com</a>	National Univ Sci and Technology, Islamabad, Pakistan	Oral
069	BRM	CLINICAL & RESEARCH LABORATORIES IN PESHAWAR CITY OF KP PROVINCE: ANALYSIS OF BIOSECURITY MEASURES PREDICT INSECURITY <a href="mailto:hoorshumail.bioinformatics@kmu.ed.pk">hoorshumail.bioinformatics@kmu.ed.pk</a>	Khyber Medical University, Peshawar, Pakistan	Oral
070	AP	ECOLOGICAL BASED FEED INVENTIONS FOR LAMBS/KIDS FATTENING MANEUVER TO EARN GREATER PROFIT REVENUES DURING EID-UL-AZHA MARKETING IN PAKISTAN Abdul Razzaq*; <a href="mailto:abdulrazzaqazrc@gmail.com">abdulrazzaqazrc@gmail.com</a>	Agriculture Research in the Dry Areas (ICARDA), Jordan	Oral
071	AP	IDENTIFICATION OF NON-GENETIC FACTORS AFFECTING WEIGHTS AT BIRTH, PRE-POST WEANING, AND GREASY FLEECE PERFORMANCE OF KAJLI SHEEP IN PAKISTAN Abdul Razzaq*; <a href="mailto:abdulrazzaqazrc@gmail.com">abdulrazzaqazrc@gmail.com</a>	Agriculture Research in the Dry Areas (ICARDA), Jordan	Poster

Code	Category	Title/authors/corresponding email	Institution	Presentation
072	AH	<p>EPIDEMIOLOGICAL RISK FACTORS, PROPHYLACTIC AND TREATMENT ARRAYS AGAINST PESTE DES PETITS RUMINANTS (PPR) IN DG KHAN, PAKISTAN</p> <p>Mahboob Ali, M Avais, Jawaria Ali Khan, A Razzaq, Farman Ullah, Zia ur Rehman*</p> <p><a href="mailto:drzia80@aup.edu.pk">drzia80@aup.edu.pk</a></p>	<p>Key Laboratory of Agricultural Animal Genetics, Breeding and Reproduction, Education, Huazhong Agricultural University, Wuhan 430070, Peoples Republic of China</p>	
073	ED	<p>DETERMINANTS OF LABOR FORCE PARTICIPATION OF LITERATE FEMALES – SCOPE FOR CONTRIBUTION IN FOOD VALUE CHAIN</p> <p>Samina Naz Qureshi, Rehmat Ali, Sadia Subhan Qureshi, Shah Murad Khan, Mamoon Ahmed and Irfan ul Haq Qureshi* <a href="mailto:i.quraysh@gmail.com">i.quraysh@gmail.com</a></p>	<p>Institute of Management Sciences, Peshawar-25000, Pakistan/Dairy Science Park, Peshawar</p>	
074	AP	<p>HOMOFERMENTATIVE BACTERIAL INOCULATION, NUTRITIVE QUALITY AND ASPERGILLUS TOXINS EVALUATION OF ALFALFA SILAGE MEDICAGO SATIVA L.</p> <p>Nafeesa Qudsia Hanif*, Sojida Latif, Anum Iftikhar, Khush Bakht, Iffat Tohiro, Nighat Sultana, Kishwar Sultana and Mujahid Hassan</p> <p><a href="mailto:Nafeesahanif9@gmail.com">Nafeesahanif9@gmail.com</a></p>	<p>AgriPak Labs, Pakistan - NTI (Licensee Romer Labs, Inc., USA)</p>	
075	PH	<p>OCCURRENCE OF MYCOBACTERIUM BOVIS IN BOVINE MILK AND CONSUMERS' KNOWLEDGE, ATTITUDE AND PRACTICES TOWARD TUBERCULOSIS IN PESHAWAR, PAKISTAN.</p> <p>Irfan Khattak*, M Hassan Mushtaq, Sultan Ayaz, Sajid Ali, Sadeeq Ur Rahman, Javed Muhammad, Haq Amanullah, M Luqman Sohail, Furqan Awan, Tahir Usman</p> <p><a href="mailto:irfankhattak@awkum.edu.pk">irfankhattak@awkum.edu.pk</a></p>	<p>College of Veterinary Sciences &amp; Animal Husbandry, Abdul Wali Khan University Mardan, Pakistan</p>	

Code	Category	Title/authors/corresponding email	Institution	Presentation
076	AP	EFFECT OF YEAST CULTURE (SACCHROMYCES CEREVISIAE) ON LACTOGENESIS IN ACHAI CATTLE Muhammad Hanif Khan, Syed M Suhail*, M Subhan Qureshi, Hayaz uddin, Ijaz Ahmed, Iftikhar Ahmed and Khalid Khan <a href="mailto:syedmuhammadsuhail@yahoo.com">syedmuhammadsuhail@yahoo.com</a>	Department of Livestock Management and Animal Breeding and Genetics, University of Agriculture Peshawar-Pakistan	
077	AP	STUDY ON GROWTH PARAMETERS AND TRANSMISSION OF GENETIC WORTH IN LOCAL RABBITS Khalid Khan, Syed Muhammad Suhail*, Rifatullah Khan, Sarzamin Khan, Muhammad Subhan Qureshi, and Iftikhar Ahmad <a href="mailto:syedmuhammadsuhail@yahoo.com">syedmuhammadsuhail@yahoo.com</a>	Faculty of Animal Husbandry and Veterinary Sciences, University of Agriculture, Peshawar	
078	AP	IMPACT OF GRADED LEVELS OF INOCULANT ON MICROBIOLOGICAL AND NUTRITIONAL QUALITY OF CORN SILAGE Smayia Sharif, Shakira Ghazanfar, Shahid Zaman, Iffat Tahira, Nafeesa Qudsia Hanif and M Imran* <a href="mailto:m_imran766@hotmail.com">m_imran766@hotmail.com</a>	Department of Microbiology, Quaid-i-Azam University, Islamabad-45320, Pakistan	
079	BRM	INVESTIGATING DISINFECTION BY-PRODUCTS FORMATION POTENTIAL OF CHLORINE AND CHLORAMINE IN DRINKING WATER Amrah Qureshi, Imran Hashmi*, Romana Khan <a href="mailto:imranhashmi@iese.nust.edu.pk">imranhashmi@iese.nust.edu.pk</a>	National University of Sciences and Technology, Islamabad, Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
080	AP	EFFECT OF REPLACEMENT OF MAIZE FODDER WITH MAIZE SILAGE IN TOTAL MIXED RATION ON NUTRIENT INTAKE, DIGESTIBILITY AND PRODUCTION PERFORMANCE OF EARLY LACTATING NILI-RAVI BUFFALOES Rafi-uddin*, Muhammad Abdullah, Khalid Javed, , Ghulam Hussain Jaffar, Muhammad Farooq Tareen, Muhammad Akbar Nasar <a href="mailto:rafi_kaka27@yahoo.com">rafi_kaka27@yahoo.com</a>	Livestock and Dairy Development Department Balochistan	
081	MT	SOME TECHNOLOGICAL PROPERTIES OF MEAT PROTEIN Halime Alp*, Mustafa Karakaya Halime Alp <a href="mailto:halimealp@hotmail.com">halimealp@hotmail.com</a>	Selcuk University KarapınarAydoğanlar Vocational School, Food Processing Department, Konya / Turkey	
082	BRM	FROM DAIRY SCIENCE PARK TO 'ONE HEALTH' SECURITY: ACHIEVEMENTS; CHALLENGES FOR GOVERNMENT & ACADEMIA IN KHYBER PUKHTUNKHWA, PAKISTAN. Tayyab ur Rehman* <a href="mailto:tayyab.ibms@kmu.edu.pk">tayyab.ibms@kmu.edu.pk</a>	Khyber Medical University Peshawar	
083	DT	INDOOR POSITIONING AND NAVIGATION USING BEACONS TECHNOLOGY Samia Subhan Qureshi*, Mehak Asad, Sidra Sayed, Mohammad Haseeb Zafar * <a href="mailto:samiasq@gmail.com">samiasq@gmail.com</a>	University of Engineering and Technology, Peshawar	
084	PH	ANTIMICROBIAL PATTERN COMBINED ACTION EFFECT BASED TOOTHPASTE AND PLANT EXTRACT ON ORAL ISOLATES Muna Jalal Ali*, Essam A. Makky <a href="mailto:munajalal2@gmail.com">munajalal2@gmail.com</a>	Universiti Malaysia Pahang, Malaysia	



Code	Category	Title/authors/corresponding email	Institution	Presentation
085	AP	DETERMINATION OF BREEDING OBJECTIVES AND ECONOMIC WEIGHTS IN ZANDI SHEEP REARED AT SEMI-INTENSIVE REARING SYSTEM OF QOM PROVINCE IN IRAN MAHDI KHOJASTEHKEYI, MAJID KALANTAR Seyed Mahdi Hosseini*, Farmanullah, Zia ur Rehman, Sima Savar Sofla, Li Guo Yang <a href="mailto:mehdi_hoseini10@yahoo.com">mehdi_hoseini10@yahoo.com</a>	Animal Science Department, Qom Agricultural and Natural Research and Education Center, AREEO, Qom, Iran/ Key Laboratory of Agricultural Animal Genetics, Breeding and Reproduction, Education Ministry of China, College of Animal Science and Thechnology, Huazhong Agricultural University, Wuhan, Hubei 430070, China	
086	AP	BIOFERTILIZERS FOR SUSTAINABLE AGRICULTURE AND FOOD SECURITY IN PAKISTAN Zahir Shah* and Tasneem Shah <a href="mailto:zahirshah@aup.edu.pk">zahirshah@aup.edu.pk</a>	Department of Soil & Environmental Sciences The University of Agriculture, Peshawar	
087	PH	CRYPTOSPORIDIUM: AN EMERGING ZONOSIS IN SOUTHERN KHYBER PAKHTUNKHWA (KPK), PAKISTAN Muhammad Luqman Sohail, Naimat Ullah Khan, Muhammad Hassan Saleem, Aneela Zameer Durrani, Nisar Ahmad, Ayesha Hassan, Sultan Ayaz <a href="mailto:Vet.lugman@gmail.com">Vet.lugman@gmail.com</a>	University College of veterinary and Animal Sciences, The Islamia University of Bahawalpur, Pakistan	
088	AP	IMPACTS OF USING IMPROVED SEED OF BERSEEM CLOVER ( <i>TRIFOLIUM ALEXANDRINUM</i> L.) ON FORAGE QUALITY, YIELDS AND NET INCOME OF SMALLHOLDER DAIRY FARMERS Muhammad Shoaib Tufail* <a href="mailto:mtufail@csu.edu.au">mtufail@csu.edu.au</a>	Graham Centre for Agricultural Innovations, Charles Sturt University, Wagga Wagga 2678, NSW, Australia	

Code	Category	Title/authors/corresponding email	Institution	Presentation
089	AH	DIAGNOSIS OF LATENT BACTERIAL INFECTIONS THROUGH ACUTE PHASE PROTEINS (APPs) IN BOVINES Shafia Tehseen Gul, Munazzah Fatima, Muhammad Kashif Saleemi, Ahrar Khan, Maqbool Ahmad and Muhammad Kasib Khan Shafia Tehseen Gul* <a href="mailto:drshafia66@yahoo.com">drshafia66@yahoo.com</a>	University of Agriculture, Faisalabad, Pakistan	
090	AP	ANIMAL BASED ORGANIC CARBON SOURCES ENHANCE CROP PRODUCTIVITY AND SOIL SUSTAINABILITY: A GOOD STRATEGY TO COMBAT FOOD SECURITY ISSUE IN DEVELOPING COUNTRIES Amanullah Jan* Email: <a href="mailto:amanullah@aup.edu.pk">amanullah@aup.edu.pk</a>	Faculty of Crop Production Science, The University of Agriculture, Peshawar- 25130	
091	MT	EFFECTS OF OLIVE LEAVES ON SHELF LIFE OF BEEF PATTIES Gamze Acar*, Mustafa Karakaya, Ali Samet Babaoğlu, Kübra Ünal <a href="mailto:gamze.kolay1@gmail.com">gamze.kolay1@gmail.com</a>	Agriculture Faculty, Food Engineering Department, Selcuk University, Konya, Turkey	
092	AH	TWO NOVEL SNPS IN STAT3 GENE ARE ASSOCIATED WITH MASTITIS SUSCEPTIBILITY IN DAIRY CATTLE Tahir Usman*, Sultan Ayaz, Irfan Khattak, Naimat Ullah Khan, Saira Gul and Atta ur Rehman <a href="mailto:tahircau@gmail.com">tahircau@gmail.com</a>	College of Veterinary Sciences and Animal Husbandry, Abdul Wali Khan University Mardan 23200, Pakistan	
093	PH	OCCURANCE AND CAUSES OF INSOMNIA IN YOUTH OF BAHAWALPUR DIVISION OF PAKISTAN M Khurram Waqas, Anum Saqib <a href="mailto:mkhurramwaqas@gmail.com">mkhurramwaqas@gmail.com</a>	University of Veterinary and Animal Sciences, Lahore, Pakistan	
094	AH	GENOTOXIC AND TOXICOPATHOLOGICAL EFFECT OF AFLATOXIN B <sub>1</sub> IN GRASS CARP ( <i>Ctenopharyngodon idella</i> ) Hayatullah Khan, Umer Saddique* <a href="mailto:druskhattak@yahoo.com">druskhattak@yahoo.com</a>	Department of Animal Health, The University of Agriculture Peshawar-Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
095	AH	IN-VITRO CYTOTOXIC AND ANTIVIRAL ACTIVITIES OF EXTRACTS OF EQUISETUM DEBILE AGAINST FOOT AND MOUTH DISEASE VIRUS Naeem Rasool, Muhammad Ovais Omer*, M Ashraf, Imran Altaf <a href="mailto:ovomer@gmail.com">ovomer@gmail.com</a>	Department of Pharmacology and Toxicology, University of Veterinary and Animal Sciences, Lahore, Pakistan.	
096	BRM	CRIMEAN CONGO HEMORRHAGIC FEVER (CCHF) IN THE SCENARIO OF DSP-BIO-RISK MANAGEMENT IN PAKISTAN Parkha Riaz Nasrullah, M.A. Kakar*, Azizullah, M.S. Qureshi and Mithat Direk <a href="mailto:adagul@yahoo.com">adagul@yahoo.com</a>	Department of Human Nutrition, University of Agriculture, Khyber Pakhtunkhwa, Pakistan	
097	AP	TOTAL TRACT NUTRIENT RETENTION AND APPARENT METABOLIZABLE ENERGY OF DIFFERENT CEREALS AND THEIR VARIETIES BY QUAILS Muhammad Sohail, Asad Sultan, Sarzamin Khan, Naila Imtiaz, Sher Bahadar Khan* <a href="mailto:asadzia2003@yahoo.com">asadzia2003@yahoo.com</a>	Department of Poultry Science, * Animal Health, The University of Agriculture, Peshawar, Pakistan Dr. Asad Sultan	
098	AH	POINT PREVALENCE OF GASTROINTESTINAL HELMINTHES IN CATTLE AND BUFFALO IN DISTRICT KASUR, PAKISTAN Inam Ullah, M Hassan Saleem, Syed Saleem Ahmad, M Haroon Akbar, Farmanullah, A Razzaq, Zahida Fatima, Zia Ur Rehman, M Salim <a href="mailto:farman_aup@yahoo.com">farman_aup@yahoo.com</a>	Department of Animal Breeding and Genetics, Lasbela University of Agriculture, Water and Marine Sciences, Uthal, Balochistan, Pakistan	
099	AP	IMPACT OF GRADED LEVELS OF INOCULANT ON MICROBIOLOGICAL AND NUTRITIONAL QUALITY OF CORN SILAGE Smayia Sharif, Shakira Ghazanfar, Shahid Zaman, Iffat Tahira Nafeesa Qudsia Hanif and M Imran* <a href="mailto:m_imran766@hotmail.com">m_imran766@hotmail.com</a>	Department of Microbiology, Faculty of Biological Sciences, Quaid-i-Azam University, Islamabad-45320, Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
100	AP	EFFECT OF SACCHAROMYCES CEREVISIAE INCORPORATION AS PARTIAL FOR REPLACEMENT FOR SOYBEAN MEAL ON OVERALL PERFORMANCE OF BROILERS AT FINISHER PHASE M Inam, Sarzamin Khan, Asad Sultan, Nazir A Khan, M Subhan Qureshi <a href="mailto:dr.inam@sbbu.edu.pk">dr.inam@sbbu.edu.pk</a>	Shaheed Benazir Bhutto University Sheringal Dir Upper; KPK-Pakistan	
101	AP	THE SIGNIFICANCE OF SELENIUM, COBALT AND MAGNESIUM, AND THE EFFECTS OF ITS DEFICIENCY IN ANIMAL HEALTH Abdul Malik Tareen, Kaleemullah, Habib ur Rehman*, Amir Rasool, Ihsanullah kakar, M Azam kakar, Saeed ur Rehman, Kamran taj, Tauseef M Asmat, Ahmad K Tareen Habib-Ur-Rehman* <a href="mailto:hur_2085qta@yahoo.com">hur_2085qta@yahoo.com</a>	Department of Microbiology, University of Balochistan (UoB), Quetta, Pakistan	
102	FSE	A STUDY OF STRUCTURAL CHANGES IN LIVESTOCK ECONOMY OF PAKISTAN: AN EMPIRICAL ANALYSIS Hafiz Zahid Mahmood, Asghar Ali, Sarfraz Hassan and Khalid Mushtaq <a href="mailto:asghar.ali@uaf.edu.pk">asghar.ali@uaf.edu.pk</a>	Institute of Agricultural and Resource Economics, University of Agriculture, Faisalabad, Pakistan	
103	AH	INCIDENCE RATE OF BLOOD PROTOZOANS IN DAIRY CATTLE IN KOHAT AND SURROUNDINGS M Rashid Khan, Zabita Khan and Shams-ul-Hayat <a href="mailto:shamsulhayat55@gmail.com">shamsulhayat55@gmail.com</a>	Veterinary Research and Disease Investigation Centre, Hangu Road Kohat	

Code	Category	Title/authors/corresponding email	Institution	Presentation
104	AP	UPSHOT OF INHERITED FACTORS ON PROLIFIC AND REPRODUCTIVE RECITAL TRAITS OF KAJLI SHEEP IN PAKISTAN Farmanullah*, Khalid Javed, Mohammad Salim, Momen Khan, Sajid Ali, Abdul Razzaq, Ahmad Nawaz Khosa, Muhammad Altaf Hussain, Sibghatullah, Ziaur Rahman, Dinesh Bhattiarai, Talpur Hira Sajjad <a href="mailto:farman_aup@yahoo.com">farman_aup@yahoo.com</a>	Key Laboratory of Agricultural Animal Genetics, Breeding and Reproduction, Education Ministry of China, College of Animal Science and Technology, Huazhong Agricultural University, Wuhan 430070, People's Republic of China	
105	ED	MONOPOLISTIC COMPETITION AND PRODUCT RIVALRY: AN ANALYSIS OF PROFIT MAXIMIZATION IN DAIRY FOOD PRODUCTS Ghazala Yasmeen*, Maleeha Khan <a href="mailto:Ghazala_nizam2006@yahoo.com">Ghazala_nizam2006@yahoo.com</a>	Women University Mardan, Pakistan.	
106	AH	IMPACT OF CLIMATE CHANGE ON EPIDEMIOLOGY OF TICK-BORNE HAEMOPARASITES IN AZAKHELI BUFFALO IN DISTRICT SWAT, PAKISTAN Zahida Fatima*, Amjad Khan, M Hassan Mushtaq, Mansur ud Din Ahmad, Asghar Khan, Shahid Hussain Farooqi, Abdul Razzaq, Aamanullah, M Luqman Sohail <a href="mailto:zahidavet@yahoo.com">zahidavet@yahoo.com</a>	Pakistan Agricultural Research Council, Islamabad, Pakistan	
107	FSE	SODIUM SILICATE SEED PRIMING LEADS TO ENHANCED ANTIOXIDATIVE PROTECTION OF WHEAT ( <i>TRITICUMAESTIVUM L.</i> ) SEEDLINGS UNDER WATER-DEFICIT STRESS Arruje Hameed*, Tahir Farooq, Amjad Hameed, Sadia Javed, M Ibrahim, Munir Ahmad Sheikh <a href="mailto:arrujetahirfsd@gmail.com">arrujetahirfsd@gmail.com</a>	Department of Biochemistry, Government College University, Faisalabad- Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
108	AH	DIETARY BUTTER INDUCES METABOLIC DISORDER IN RAT MODEL Maha Saleem *, M Naeem Faisal, Haseeb Anwar, Ahmed Abdulsadiq, Aisha Mehmood, Humaira Muzaffar, Hira Javed, Farooq Azam, Tanweer Khaliq <a href="mailto:dr.hirawarraich443@gmail.com">dr.hirawarraich443@gmail.com</a>	Institute of Pharmacy, Physiology & Pharmacology, University of Agriculture, Faisalabad, Pakistan	
109	AH	CURRENT QUINOLONES SHOW SIGNIFICANT HYPOGLYCEMIC EFFECTS IN EUGLYCEMIC RABBITS Niaz Ali, Zia Ud Din, M Sajid, Sarzamin Khan, M Subhan Qureshi, Shakir Ullah, M Nabi, Waqar Ali <a href="mailto:niazpharmacist@yahoo.com">niazpharmacist@yahoo.com</a>	Department of Pharmacology, Institute of Medical Sciences, Khyber Medical University, Kohat, Khyber Pakhtunkhwa, Pakistan.	
110	FS	BENZIMIDAZOLE AND THIOBENZIMIDAZOLE INDUCED CHANGES IN GROWTH AND ANTIOXIDANT ATTRIBUTES OF OKRA GROWN UNDER CHROMIUM STRESS Tahir Farooq,* Arruje Hameed, Azeem Sultan, Matloob Ahmed, M Sabir and Kiran Sana <a href="mailto:tahirfarooqfsd@gmail.com">tahirfarooqfsd@gmail.com</a>	Government College University, Faisalabad	
111	FSE	BALUCHISTAN NARI MASTER AND ITS IMPORTANCE FOR THE FUTURE BEEF INDUSTRY OF PAKISTAN M. Azam Kakar, Sana Durrani, Shakoor A. Baloch, Azizullah, M.Subhan Qureshi, Parkha Riaz Nasrullah and Mithat Direk <a href="mailto:adagul@yahoo.com">adagul@yahoo.com</a>	The World Bank Pakistan, 18-B, Jinnah Town, Samungli Road, Quetta, Pakistan	
112	FSE	ROLE OF BIOTECHNOLOGY IN THE PROSPERITY AND FOOD SECURITY OF DEVELOPING WORLD M. Azam Kakar, Sana Durrani, Shakoor A. Baloch, Azizullah, M Subhan Qureshi, Parkha Riaz Nasrullah and Mithat Direk <a href="mailto:adagul@yahoo.com">adagul@yahoo.com</a>	The World Bank Pakistan, 18-B, Jinnah Town, Samungli Road, Quetta, Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
113	ED	ROLE OF WOMEN IN LIVESTOCK AND DAIRY DEVELOPMENT IN BALOCHISTAN, PAKISTAN Sana Durrani*, Shakoor A. Baloch, Parkha Riaz Nasrullah, M.A. Kakar, Azizullah, M Subhan Qureshi and Mithat Direk * <a href="mailto:ceo.two@gmail.com">ceo.two@gmail.com</a>	Small and Medium Enterprise Development Authority, Balochistan Quetta, Pakistan	
114	FSE	ENHANCING OUTPUT OF ANIMAL-ORIGIN PRODUCTS FOR ENSURING FOOD SECURITY IN PAKISTAN; A REVIEW ARTICLE Zahida Fatima*, Abdul Razzaq, Shahid Rafique, Aman Ullah, Amjad Khan, Ayesha Tahir, Rehana Kausar <a href="mailto:zahidavet@yahoo.com">zahidavet@yahoo.com</a>	Deputy Director, Animal Sciences Division, National Agricultural Research Council, Islamabad, Pakistan	
115	PH	SEROPREVALENCE AND RISK FACTOR ANALYSIS OF HUMAN LEPTOSPIROSIS IN DISTINCT CLIMATIC REGIONS OF PAKISTAN M Luqman Sohail*, M Sarwar Khan, M Ijaz, Sajid Umar, Zahida Fatima, Omer Naseer <a href="mailto:vet.luqman@gmail.com">vet.luqman@gmail.com</a>	University College of veterinary and Animal Sciences, The Islamia University of Bahawalpur, Pakistan	Oral?
116	FSE	TOTAL FACTOR PRODUCTIVITY GROWTH OF LIVESTOCK SECTOR IN PAKISTAN: A TIME SERIES ANALYSIS Asghar Ali*, M Ikhtlaq Mansoor and Abdul Ghafoor <a href="mailto:asghar.ali@uaf.edu.pk">asghar.ali@uaf.edu.pk</a>	Institute of Agricultural and Resource Economics, University of Agriculture, Faisalabad, Pakistan	Oral?
117	AH	HIGH PROTEIN DIET LOWERS BLOOD GLUCOSE LEVEL THROUGH REGENERATION OF PANCREATIC B CELLS IN DIABETIC RAT MODEL Saima Sattar, MN Faisal, Haseeb Anwar, A Abdulsadiq, Aisha Mahmood, Humaira Muzaffar, Farooq Ahmed*, A Basit, Farooq Azam <a href="mailto:Khanfarooq391@gmail.com">Khanfarooq391@gmail.com</a>	Institute of Pharmacy, Physiology & Pharmacology, University of Agriculture, Faisalabad, Pakistan.	



Code	Category	Title/authors/corresponding email	Institution	Presentation
118	AH	COMPARATIVE EFFICACY OF LOCAL AND PARENTAL ANTIBIOTICS FOR THE TREATMENT OF CLINICAL MASTITIS IN DAIRY BUFFALOES Tanveer Ahmad, Ali Khera, Ejaz Ahmad, MS Akhtar and Asim Faraz <a href="mailto:tanveeruaf@gmail.com">tanveeruaf@gmail.com</a>	Department of Clinical Science, FVS, BZU, Multan	
119	FSE	NEED OF HALAAL CERTIFICATION FOR EDIBLES, BEVERAGES, MEDICINES, COSMETICS(AN OVERVIEW) Mufti Syed Naeem Badshah, Sadia Subhan Qureshi and Shah Murad Khan <a href="mailto:ssqureshi.arabic@gmail.com">ssqureshi.arabic@gmail.com</a>	Islamia College Peshawar	
120	PH	ANTHELMINTIC DRUGS IN THE DARK ERA OF ANTHELMINTIC RESISTANCE: REPORT FROM PAKISTAN Zia ud Din Sindhu, Rehman Hafeez, Zafar Iqbal, M Kasib Khan, RaoZahid Abbas, Bilal Aslam, M Nisar Khan <a href="mailto:ziasandhu@hotmail.com">ziasandhu@hotmail.com</a>	Department of Parasitology University of Agriculture Faisalabad	
121	AP	USE OF MOS IN BROILER DIETS: A REVIEW M.F.A.Chacher, Z. Kamran, U. Ahsan, S. Ahmad, H.G. Qutab ud dinand ö. Cengi M Furqan Asghar Chacher <a href="mailto:vetchachar@gmail.com">vetchachar@gmail.com</a>	Department of Obstetrics and Gynecology, Faculty of Veterinary Medicine, Ataturk University, 25240 Erzurum, Turkey	
122	BRM	SERO-PREVALENCE OF ECHINOCOCCOSIS (ECHINOCOCCUS GRANULOSUS) IN CAMEL POPULATION OF SELECTED ECOLOGICAL ZONE OF BALUCHISTAN, PAKISTAN M Shafi Hasni, M Imran*, M Arsahd, M Saqib, M Sohail Sajid, M Kasib Khan, Abdul Qudoos <a href="mailto:imran.asghar@uaf.edu.pk">imran.asghar@uaf.edu.pk</a>	Department of Parasitology, University of Agriculture-38040, Faisalabad, Pakistan	Accepted Oral



Code	Category	Title/authors/corresponding email	Institution	Presentation
123	AH	FREQUENCY DISTRIBUTION OF TICK (ACARI: IXODIDAE) INFESTING THE LOCAL AND EXOTIC CATTLE BREEDS OF DISTRICT SWAT, KHYBER PAKHTUNKHWA, PAKISTAN Tanveer Iqbal, M Imran*, M Nisar Khan, M Sohail Sajid, Abdul Qudoos, Asim Shamim <a href="mailto:imran.asghar@uaf.edu.pk">imran.asghar@uaf.edu.pk</a>	Department of Parasitology, University of Agriculture-38040, Faisalabad, Pakistan	Accepted Oral
124	ED	DIETARY SUPPLEMENTATION OF PROBIOTIC YEAST IMPROVES GASTROINTESTINAL TRACT MICROBITA IN GROWING SAHIWAL HEIFERS Shakira Ghazanfar,*, Iftikhar Ahmed, M Iqbal Anjum <a href="mailto:shakira_akmal@yahoo.com">shakira_akmal@yahoo.com</a>	Institute of Microbial Culture Collection, National Agricultural Research Centre, Park Road, Islamabad-45500, Pakistan	Accepted Oral
125	BRM	EPIDEMIOLOGY OF ZONOTIC BRUCELLOSIS IN LIVESTOCK FARMS IN PAKISTAN; “AN UNDERESTIMATED PUBLIC HEALTH PROBLEM” Amjad Khan, M Hassan Mushtaq, Habib un Nabi, Abdul Rehman, Mansur-Ud Din Ahmad, Shahid Hussain Farooqi and Asghar Khan. <a href="mailto:dramjadjkhan77@gmail.com">dramjadjkhan77@gmail.com</a>	University of Veterinary and Animal Sciences, Lahore, Pakistan	Accepted Oral
126	FSE	IMPROVING ABIOTIC STRESS TOLERANCE IN CEREALS: AN INTEGRATED APPROACH FOR GLOBAL FOOD SECURITY Amjad Hameed <a href="mailto:amjad46pk@yahoo.com">amjad46pk@yahoo.com</a>	Nuclear Institute for Agriculture and Biology (NIAB), Faisalabad, Pakistan	Oral, accepted
127	BRM	ANTIBIOTIC RESISTANCE AND RESISTANT GENES IN E. COLI BELONG TO DIFFERENT PHYLOGENETIC GROUPS SB Khan*, Zou Geng, Sher Ali, Tariq Shah, HA Ullah, Muqadar Shah, Q Ullah, Murad Ali, I Ahmad, Mumtaz Ali, Zhou Rui <a href="mailto:sbvvetdr@yahoo.com">sbvetdr@yahoo.com</a>	Department of Animal Health, The University of Agriculture, Peshawar, Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
128	FSE	IN VITRO ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY OF DICHLORO-METHANE EXTRACT OF RANUNCULUS MURICATUS AND CASSIA FISTULA Farooq Azam*, Bashir Ahmad, M Ashraf, Faqir M, M Naeem Faisal, Wafa Majeed, Aisha Mahmood, Humaira Muzaffar, Jawad Aslam <a href="mailto:farooq.leo@hotmail.com">farooq.leo@hotmail.com</a>	Institute of Pharmacy, Physiology and Pharmacology, University of Agriculture, Faisalabad, Pakistan	
129	AH	THE ECONOMIC LOSSES AND BURDEN OF ENDOPARASITES IN BUFFALO AND CATTLES OF RURAL PUNJAB “AN UNDERESTIMATED EMERGING THREAT” M Hassan Mushtaq*, Ghazala Nawaz, M Nawaz Malik, Amjad K <a href="mailto:hassan.mushtaq@uvas.edu.pk">hassan.mushtaq@uvas.edu.pk</a>	University of Veterinary and Animal Sciences Lahore	
130	AH	SERO-EPIDEMIOLOGY OF TOXOPLASMOSIS IN PASTORAL GOAT HERDS AND SHEPHERDS IN PUNJAB, PAKISTAN M Mushtaq*, Shams-ur-Rehman, Mansur ud Din A, A Khan <a href="mailto:hassan.mushtaq@uvas.edu.pk">hassan.mushtaq@uvas.edu.pk</a>	University of Veterinary and Animal Sciences Lahore.	
131	EP	CHALLENGES AND FUTURE SCOPE OF SME SECTOR: SPECIALLY ACCESS TO FINANCE IN BALOCHISTAN Shakoor Ahmed <a href="mailto:shakoor@smeda.org.pk">shakoor@smeda.org.pk</a>	Small & Medium Enterprises Development Authority , 15-A, Airport Road, Quetta. Pakistan	
132	AP	STUDIES ON BIOCHEMICAL ASPECTS OF SAHIWAL AND CROSS-BRED (SAHIWAL X FRIESIAN) BULLS SEMEN M Jamil Ahmad, FarmanUllah, Hafiz Ishfaq Ahmad, Zia Ur Rehman* <a href="mailto:drzia80@aup.edu.pk">drzia80@aup.edu.pk</a>	Key Laboratory of Agricultural Animal Genetics, Breeding and Reproduction, Huazhong Agricultural University, China.	

Code	Category	Title/authors/corresponding email	Institution	Presentation
133	AP	EFFECT OF FEEDING DIFFERENT LEVELS OF UREA TREATED SUGAR BEET PULP VERSUS UNTREATED SUGAR BEET PULP ON WEIGHT GAIN OF BULKHI SHEEP UNDER THE SUBTROPICAL REGION OF DISTRICT DERA ISMAIL KHAN Shoaib Sultan and Saqib Ali Rustham <a href="mailto:shoaibnutr@gmail.com">shoaibnutr@gmail.com</a>	Department of Animal Nutrition, The University of Agriculture Peshawar. Pakistan	
134	AP	STUDY PATTERN OF ASSORTATIVE MATING IN LOCAL RABBITS Inamullah, S M Suhail, Sarzamin Khan, M. Subhan Qureshi and Khalid Khan. <a href="mailto:syedmuhammadsuhail@yahoo.com">syedmuhammadsuhail@yahoo.com</a>	Faculty of AH&VS The University of Agriculture Peshawar	
135	PH	INCIDENCE OF DRUG RESIDUES IN MILK AND MEAT AT DISTRICT KARAK, KP, PAKISTAN Safat Ullah, Shakoor Ahmad*, Umersadique, Hamayun Khan, Sher Bahadur, Zahir Shah, Murad Ali Khan and Ijaz Ahmad <a href="mailto:shakoor.ahmad@aup.edu.pk">shakoor.ahmad@aup.edu.pk</a>	Department of Animal Health, The University of Agriculture Peshawar, Pakistan	
136	FSE	PAKISTAN WATER RESOURCES: DEVELOPMENT AND CONSERVATION NEEDS FOR SUSTAINABLE CROP AND LIVESTOCK PRODUCTION M Abdullah, MA Shakir*, M Naveed <a href="mailto:masoodshakir7@gmail.com">masoodshakir7@gmail.com</a>	Ayub Agricultural Research Institute, Faisalabad, Pakistan	
137	PH	CONTRIBUTION TO THE STUDY OF SOME LIPID AND LIPOPROTEIN PARAMETERS OF THE ORGANS IN OBESE RATS SUPPLEMENTED WITH GREEN MICROALGAE. Benyoub N*, Saidani F, Baba Ahmed Fz Email: <a href="mailto:benyoubnoredine@yahoo.fr">benyoubnoredine@yahoo.fr</a>	Laboratory of physiology, physiopathology, and biochemistry of nutrition University of Tlemcen Algeria.	

Code	Category	Title/authors/corresponding email	Institution	Presentation
138	FSE	THE CHANGE OF CLIMATE IN MAIZE PRODUCTION IN KHYBER PAKHTUNKHWA, PAKISTAN Ansa Javed <a href="mailto:ansaj165@gmail.com">ansaj165@gmail.com</a>	Planning and Dev Department, Women University Mardan	
139	AP	ROLE OF FEED ADDED ANISEED AS IMMUNUMODULANT AND GROWTH PROMOTER IN BROILER CHICKS Riaz Ahmad, FarmanUllah*, Abdul Razzaq, Islam zeb Khan, M Salim <a href="mailto:farman_aup@yahoo.com">farman_aup@yahoo.com</a>	Faculty of Vet and AS, Lasbela University of Agriculture, Water and Marine Sciences, Uthal, Balochistan, Pakistan	
140	AH	VACCINE; A RATIONAL CONTROL STRATEGY FOR MYCOPLASMA BOVIS INFECTION IN CATTLE Farhan Anwar Khan, Xi Chen, Ijaz Ahmad, Sher Bahadar Khan, Zahir Shah, Zia ur Rehman, Shakoor Ahmad, Umar Sadique, Aizhen Guo <a href="mailto:drfarhananwar2003@yahoo.com">drfarhananwar2003@yahoo.com</a>	Department of Animal Health, The University of Agriculture, Peshawar, Khyber Pakhtunkhwa, 25120, Pakistan	
141	AH	HOMOLOGY MODELLING AND MOLECULAR DOCKING STUDY OF CORTICOTROPHIN RELEASING HORMONE: A SUCCESSFUL APPROACH TO TREAT THE STRESS RELATED DISEASES. Shahid Ullah <a href="mailto:shahidullah@hust.edu.cn">shahidullah@hust.edu.cn</a>	College of Chemistry and Env Engg, Shenzhen University, Shenzhen, Guangdong, 518060, P. R. China	
142	FSE	PAKISTAN WATER RESOURCES: DEVELOPMENT AND CONSERVATION NEEDS FOR SUSTAINABLE CROP AND LIVESTOCK PRODUCTION Muhammad Abdullah, Masood A. Shakir and Muhammad Naveed <a href="mailto:masoodshakir7@gmail.com">masoodshakir7@gmail.com</a>	Ayub Agricultural Research Institute, Faisalabad, Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
143	BRM	MOLECULAR EPIDEMIOLOGY OF RESPIRATORY VIRUSES IN COMMERCIAL CHICKEN FLOCKS IN PAKISTAN FROM 2014 THROUGH 2016 Sajid Umar*, Angélique Teillaud, Hassan Bin Aslam, Jean-Luc Guerin1, Mariette F. Ducatez, IHAP <a href="mailto:s.umar@uaar.edu.pk">s.umar@uaar.edu.pk</a>	Université de Toulouse, INRA, ENVT, 31076 Toulouse, France	
144	BRM	AVIAN METAPNEUMOVIRUS VACCINE MODIFIES VIRUS SHEDDING AND IMMUNE RESPONSES OF AVIAN INFLUENZA (H9N2) INFECTION IN COMMERCIAL TURKEY Sajid Umar*, Qamar-un-Nisa, Abdul Ghafar, M Ali Shah, M Luqman Sohail, M Asif Yaseen. <a href="mailto:s.umar@uaar.edu.pk">s.umar@uaar.edu.pk</a>	Department of Pathobiology, PMAS Arid Agriculture university Rawalpindi (46000), Pakistan	
145	FSE	RED MEAT PRODUCTION TRENDS IN MONGOLIA Maralma Bayarsaikhan, Mithat Direk <a href="mailto:mithatdirek@gmail.com">mithatdirek@gmail.com</a>	Selçuk University, Department of Agric Economics, Konya-Turkey	
146	AH	REAL-TIME FLUORESCENCE LOOP-MEDIATED ISOTHERMAL AMPLIFICATION FOR DETECTION OF hCD39 IN TRANSGENIC PIG Jin Li, Wei Liu, Jun-chao Li, Hong-yan Jin, GangLi * <a href="mailto:*ligang03@caas.cn">*ligang03@caas.cn</a>	Institute of Animal Science, Chinese Academy of Agricultural Sciences, Beijing 100193,China	
147	DT	EFFECT OF SMOKING ON PHYSICO-CHEMICAL PROPERTIES AND SENSORY ATTRIBUTES OF MUDAFFARA CHEESE USING CORIANDRUM SATIVUM Ibtisam E. M. El Zubeir <a href="mailto:Ibtisammohamed@hotmail.com">Ibtisammohamed@hotmail.com</a>	Department of Dairy Production, Faculty of Animal Production, University of Khartoum, P. O. Box 321, Khartoum, Sudan.	

Code	Category	Title/authors/corresponding email	Institution	Presentation
148	AP	EFFECT OF GONADOTROPIN RELEASING HORMONE, HUMAN CHORIONIC GONADOTROPIN AND FLUNIXINE MEGLUMINE ON PREGNANCY RATES IN SAHIWAL COWS Muhammad Saleem Akhtar, Tanveer Ahmad, Laeeq Akbar Lodhi, Mushtaq Hussain Lashari, Ejaz Ahmad, Masood Akhtar <a href="mailto:drsaleem45@yahoo.com">drsaleem45@yahoo.com</a>	Department of Clinical Sciences, Faculty of Veterinary Sciences, Bahauddin Zakariya University, Multan, Pakistan	
149	AP	IMMUNOSTIMULANT AND HEPATOPROTECTIVE EFFECT OF POLY HERBS (WITHANIASOMNIFERA, LIQUORICE, ALLIUM SATIVUM AND BERBERISLYCIUM) MIXTURE EXTRACT IN BROILER PRODUCTION Noor ul Baseer*, F Ullah, Abdul Razzaq, MA Hussain. M Salim <a href="mailto:farman@webmail.hzau.edu.cn">farman@webmail.hzau.edu.cn</a>	Faculty of Animal Husbandry and Veterinary and Animal Sciences, AUP, Pakistan	
150	AH	THE FACTORS REGULATING MELANOGENESIS: A REVIEW Hira Sajjad Talpur*, Iqra Bano <a href="mailto:drhrasajjad@gmail.com">drhrasajjad@gmail.com</a>	Key Laboratory of Agric Animal Genetics, Huazhong Agric Univ, Wuhan-430070, People's Republic of China.	
151	AH	A CROSS SECTIONAL SURVEY ON GASTROINTESTINAL PARASITES IN BACKYARD POULTRY IN PAKISTAN MH Mushtaq, Ghazala Nawaz, MN Malik, Amjad Khan <a href="mailto:hassan.mushtaq@uvas.edu.pk">hassan.mushtaq@uvas.edu.pk</a>	Department of Epidemiology, University of Veterinary and Animal Sciences, Lahore, 54000, Pakistan	
152	FSE	PERMISSIBLE AND IMPERMISSIBLE FOOD/DRINK IN LIGHT OF MAZAHIB-E-ARBA 'A Dr Rashida Parveen <a href="mailto:drrashidaisb@gmail.com">drrashidaisb@gmail.com</a>	Women University Mardan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
153	ED	<p>A SIMPLE AND SENSITIVE HPLC METHOD FOR DETERMINATION OF CEFQUINOME IN PLASMA: APPLICATION TO PHARMACOKINETIC STUDY IN CATTLES.</p> <p>Ijaz Ahmad*, Haihong Hao, Jun Li, Pascal Sanders, Sher Bahadar Khan, Farhan Anwar Khan, Zahir Shah, Shuyu Xie, Saeed Ahmed, Lingli Huang, Zonghui Yuan  <a href="mailto:drijaz@aup.edu.pk">drijaz@aup.edu.pk</a></p>	Department of Animal Health, The University of Agriculture Peshawar, Pakistan	
154	FSE	<p>ESTIMATING THE CROP LOSSES CAUSED BY FUSARIUM WILT OF CHILI AND ITS CONTROL BY TARGETING MITOCHONDRIAL RESPIRATORY SYSTEM WITH NOVEL INHIBITORS USING COMPUTER AIDED DRUG DESIGNING</p> <p>Sehrish Iftikhar, Ahmad Ali Shahid, Kiran Nawaz, Waheed Anwar  <a href="mailto:Sehrish.iftikhar05@gmail.com">Sehrish.iftikhar05@gmail.com</a></p>	Institute of Agricultural Sciences, University of the Punjab, Lahore, Pakistan	
155	AH	<p>ISOLATION AND CHARACTERIZATION OF BACILLUS SPECIES EXHIBITING ANTIBACTERIAL ACTIVITY FROM LAKKI MARWAT, PAKISTAN</p> <p>Anum Tabassum, Dr. M Qasim, H Rehman, Nayab &amp; Asia Begum  <a href="mailto:anum.microbiologist@yahoo.com">anum.microbiologist@yahoo.com</a></p>	Women University Mardan, KPK Pakistan	
156	PH	<p>ANTIDIABITIC EFFECTS OF CAMEL MILK: SHORT AND LONG-TERM STUDIES ON MATERNAL UTERUS AND NEONATES SELECTED ORGANS</p> <p>Anas S Qureshi, MK Ateeq, M Usman, MZ Ali, Muqader Shah, WA Khamas  <a href="mailto:muqader849@aup.edu.pk">muqader849@aup.edu.pk</a></p>	Department of Anatomy, University of Agriculture, Faisalabad, Pakistan	



Code	Category	Title/authors/corresponding email	Institution	Presentation
157	AH	TOXICOPATHOLOGICAL EFFECT OF DIETRY OCHRATOXIN A IN QUAILS (COTURNIX JAPONICA) Faisal Ahmad, Umer Saddique <a href="mailto:drdfaisal.vet31@gmail.com">drdfaisal.vet31@gmail.com</a>	FAHV&S. The University of Agriculture Peshawar, Khyber Pukhtunkhwa-Pakistan	
158	BRM	COMPARATIVE DISTRIBUTION AND MORPHOLOGICAL FEATURES OF FIVE SPECIES OF THE GENUS PIPISTRELLUS OF MALAKAND DIVISION M Salim, Arshad Javid, Sajid Ali, FarmanUllah, Ahmad Hussain, Ume Habiba and Momen Khan <a href="mailto:mohammadsalim@uoh.edu.pk">mohammadsalim@uoh.edu.pk</a>	Department of Forestry and Wildlife Management, University of Haripur, Khyber Pakhtunkhwa, Pakistan	
159	AP	EFFECTS OF FORMER SIZMA (KONYA) MERCURY MINE ON MILK PRODUCTION Fatih Ahmet GÖKTÜRK <a href="mailto:fatihahmetgokturk@hotmail.com">fatihahmetgokturk@hotmail.com</a>	Selcuk University ,Environmental Engineering Department, Konya Turkey	
160	BRM	GOVERNMENT- ACADEMIA: MISSING LINKS AND TRUST DEFICIT WITH SPECIAL REFERENCE TO BIORISK MANAGEMENT IN KP PROVINCE. FarzanaNazli <a href="mailto:Farzana562@gmail.com">Farzana562@gmail.com</a>	Education consultant, Development sector KP Peshawar	
161	AH	A CLINICAL SURVEY OF VIRAL RESPIRATORY DISEASES OF COMMERCIAL POULTRY IN PAKISTAN M Luqman Sohail, Sajid Umar, M Usman <a href="mailto:Vet.luqman@gmail.com">Vet.luqman@gmail.com</a>	University College of veterinary and Animal Sciences, The Islamia University of Bahawalpur, Pakistan	
162	MT	EFFECTS OF GRAPE LEAF POWDERS DRIED BY MICROWAVE ENERGY ON SOME QUALITY PROPERTIES OF BEEF PATTIES AlimeYaldız, CemalettinSarıçoban, KübraÜnal <a href="mailto:alimeyldz-92@hotmail.com">alimeyldz-92@hotmail.com</a>	Agriculture Faculty, Food Engineering Department, Selcuk University, Konya, Turkey	



Code	Category	Title/authors/corresponding email	Institution	Presentation
163	MT	EFFECTS OF GROUND CRESS AND FLAX SEEDS ON SOME QUALITY PROPERTIES OF TAIL FAT Gülsüm Atasoy, Cemalettin Sarıçoban, Alime Yıldız <a href="mailto:alimeyldz-92@hotmail.com">alimeyldz-92@hotmail.com</a>	Agriculture Faculty, Food Engineering Department, Selçuk University, Konya, Turkey	
164	FSE	MICROBIAL POTENTIAL (METHANE PRODUCING MICROBE) ON METHANE PRODUCTION AT RICE-ROOTED AREA ON THAILAND JASMINE RICE FIELD Sansanee Chawanakul <a href="mailto:cha.sansanee@gmail.com">cha.sansanee@gmail.com</a>	Department of Biological Sciences, Faculty of Science, Ubon Ratchathani University, Warinchamrab district, Ubon Ratchathani 34190, Thailand	
165	DT	THE EFFECT OF SOYBEAN OR FLAXSEED OILS TO AFFECT FEED UTILIZATION, MILK PRODUCTION AND COMPOSITION, IN LACTATING COWS AM Kholif*, M Abdel Aziz, AF Sayed, AE Kholif, Hoda Hassan, GA Gouda, Layla Khaled, M Fawzy; <a href="mailto:am_kholif2000@yahoo.co.uk">am_kholif2000@yahoo.co.uk</a>	Dairy Science Department, National Research Centre, Dokki, Giza, Egypt	
166	DT	IMPLEMENTATION OF MODEL FOR IMPROVEMENT (PDCA-CYCLE) IN DAIRY LABORATORIES Abdelkader M Kholif, Dina S. Abou el Hassan, MA Khorshid and Elsayed A Elsherpieny <a href="mailto:am_kholif@hotmail.com">am_kholif@hotmail.com</a>	Dairy Science Department, National Research Centre, 12622, Dokki, Giza, Egypt	
167	AP	EFFECT OF SPRING MAIZE GENOTYPES HARVESTED AT DIFFERENT MATURITY STAGES ON YIELD, NUTRIENT COMPOSITION, FIBER DIGESTIBILITY AND SILAGE FERMENTATION QUALITY Nadar Khan* , Nazir A Khan, Nazir Ahmad, M.Subhan Qureshi, MNM Ibrahim; <a href="mailto:nadar73vet@gmail.com">nadar73vet@gmail.com</a>	The University of Agriculture Peshawar, Khyber Pakhtunkhwa, Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
168	AP	SCREENING OF TRADITIONAL AND NOVEL SUMMER MAIZE CULTIVARS AT DIFFERENT HARVEST MATURITY FOR SILAGE PRODUCTION IN NORTHERN AREAS OF PAKISTAN Nadar Khan, Nazir Ahmad Khan, Nazir Ahmad, M.Subhan Qureshi and M. N.M. Ibrahim <a href="mailto:nadar73vet@gmail.com">nadar73vet@gmail.com</a>	The University of Agriculture Peshawar, Khyber Pakhtunkhwa, Pakistan	
169	FSE	MILK PRODUCTION AROUND KONYA CITY CENTRE AND ITS RELATION WITH REGION ENVIRONMENT Sukru Dursun <a href="mailto:sdursun@selcuk.edu.tr">sdursun@selcuk.edu.tr</a>	Selcuk University, Environmental Engineering Department, Konya, Turkey	
170	APE	INFLUENCE OF FLAXSEEDS ON RUMINAL FERMENTATION AND MILK PRODUCTION Syeda Maryam Hussain, Valdo Herling, Paulo Henrique Mazza and Marcos Barbosa <a href="mailto:maryamhussain05@gmail.com">maryamhussain05@gmail.com</a>	Faculty of Animal Sciences and Food Engineering, University of Sao Paulo, Brazil.	
171	FSE	DAIRYSECTOR IN YEMEN – A DRIVING FORCE FOR FOOD SECURITY Hafsa AL Yamani*, Dehouegnon Jerry Agossou, Nazan Koluman <a href="mailto:m7y11@ymail.com">m7y11@ymail.com</a>	Department of Animal Science, Çukurova University, 01330, Adana Turkey	
172	FSE	TRAJECTORY OF WOMEN SEXUAL IMPEDIMENTS: A CONSTITUTIONAL PERSPECTIVE Frahhat Jabeen, Ansa Javed Khan <a href="mailto:ansaj165@gmail.com">ansaj165@gmail.com</a>	Women University Mardan	
173	FSE	POPULATION ESTIMATION AND THREATS TO INDIAN HARE (LEPUSNIGRICOLLIS) IN RAKH-SARDARAN GAME RESERVE, DISTRICT HARIPUR. Murtaza Ali, Ume Habiba*, M Salim, Ahmed Hussain, Farmanullah <a href="mailto:ume.habiba8@gmail.com">ume.habiba8@gmail.com</a>	Department of Forestry and Wildlife Management, University of Haripur, Khyber Pakhtunkhwa, Pakistan	

Code	Category	Title/authors/corresponding email	Institution	Presentation
174	DT	ESTIMATION OF MILK PROTEASE ACTIVITY FROM UNINFECTED MAMMARY GLANDS OF NILI-RAVI BUFFALOS, SAHIWAL AND CROSS-BREED COWS OF PAKISTAN. RaziaKausar*, AmjadHameed, and Zafar Iqbal Qureshi <a href="mailto:razia_uaf@hotmail.com">razia_uaf@hotmail.com</a>	Department of Anatomy, FVS, University of Agriculture, FSD Nuclear Institute of Agriculture and Biology	
175	FS	EVALUATION OF LIVESTOCK PRODUCTION IN ÇELEBILER VILLAGE WITH SPECIAL EMPHASIS ON PASTURE Sema Yaman* <a href="mailto:semayaman@ohu.edu.tr">semayaman@ohu.edu.tr</a>	Niğde Ömer Halisdemir University Faculty of Agricultural Sciences and Technologies, 51240 Niğde, Turkey	
176	FS	USE OF PROPOLIS IN RUMINANT NUTRITION Sema Yaman, Sibel Canoğulları Doğan, İlknur Uçak <a href="mailto:semayaman@ohu.edu.tr">semayaman@ohu.edu.tr</a>	Niğde Ömer Halisdemir University, Faculty Agric Sci & Tech, Niğde, Turkey	
177	FS	AN ALTERNATIVE PROTEIN SOURCE-INSECTS IN ANIMALNUTRITION Mustafa Boğa, Sema Yaman* <a href="mailto:semayaman@ohu.edu.tr">semayaman@ohu.edu.tr</a>	Niğde Ömer Halisdemir Üniversitesi Vocational School, Bor, Niğde, Turkey	
178	BRM	ANTIMICROBIAL RESISTANCE IN THE FOOD CHAIN IN DEVELOPING COUNTRIES AND PUBLIC HEALTH CONSEQUENCES: A CALL FOR ACTION Sahibzada Shafiullah, M Subhan Qureshi; <a href="mailto:sshafiullah@csu.edu.au">sshafiullah@csu.edu.au</a>	Charles Sturt University, Wagga Wagga, NSW 2678. Graham Centre for Agricultural Innovation	
179	MT	DETERMINATION OF THE EFFECTS OF THE ADDITION OF HYSSOP POWDER ON SOME QUALITY PROPERTIES OF BEEF PATTIES Şeyma Cabi Çarşi, Cemalettin Sariçoban, Alime Yaldiz <a href="mailto:seymacarsi@gmail.com">seymacarsi@gmail.com</a>	Agriculture Faculty, Food Engineering Department, Selcuk University, Konya, Turkey	

<b>Code</b>	<b>Category</b>	<b>Title/authors/corresponding email</b>	<b>Institution</b>	<b>Presentation</b>
180	BRM	FROM DAIRY SCIENCE PARK TO 'ONE HEALTH' SECURITY: CHALLENGES& EXPECTATIONS FOR GOVERNMENT-ACADEMIA-INDUSTRY IN KHYBER PAKHTUNKHWA, PAKISTAN Tayyab Rehman <a href="mailto:tayyab.ibms@kmu.edu.pk">tayyab.ibms@kmu.edu.pk</a>	Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, Pakistan	
181	FS	WHOLE GENOME SEQUENCING OF CATTLE BREEDS FROM PAKISTAN AND THEIR USE IN GENOMIC SELECTION FOR ENHANCING MILK AND MEAT PRODUCTION Shahid Mansoor <a href="mailto:shahidmansoor7@gmail.com">shahidmansoor7@gmail.com</a>	National Institute for Biotechnology and Genetic Engineering (NIBGE), P O Box, 577, Jhang Road, Faisalabad, Pakistan	

**SEROPREVALENCE AND POTENTIAL RISK FACTORS ASSOCIATED WITH BOVINE BRUCELLOSIS UNDER DIVERSE PRODUCTION SYSTEMS IN PAKISTAN**

Aman Ullah Khan, Iahasham Khan, Muhammad Younus, Syed Ehtisham-Ul-Haque, Usman Waheed, Falk Melzer, Hosny El-Adawy, Heinrich Neubauer, Muhammad Asif, Idrees

University of Veterinary & Animal Sciences, Lahore-Pakistan (Jhang Campus)

[Amanullah.khan@uvas.edu.pk](mailto:Amanullah.khan@uvas.edu.pk)

**ABSTRACT**

Brucellosis is one of the major problems affecting milk producing animals which deteriorates the health of livestock all over the world including Pakistan. It is a disease of zoonotic significance causing disease in humans. In animals economic loss is caused by infertility, orchitis, abortions and synovitis. In this study, 175 milk (90 from cattle & 85 from buffaloes) and 402 serum (208 from cattle & 194 from buffalo) samples collected from cattle and buffaloes in three selected districts (Narowal, Gujranwala and Gujrat) of the Punjab, Pakistan were investigated for anti-brucella antibodies during the months of February to June, 2015. Milk samples were tested by Milk Ring Test (MRT), while serum samples were initially screened with Rose Bengal Plate agglutination test (RBPT) and confirmed with Indirect Enzyme Linked Immunosorbant Assay (i-ELISA) by i-ELISA (IDEXX Brucellose Serum X2 Ab Test, Switzerland). In total 21.7% milk samples were positive (cattle 22.2% and buffaloes 21.1%). Brucella antibodies were detected in 16.4% of tested sera with RBPT (cattle 18.8% and buffaloes 13.9%) and 11.4% by i-ELISA (cattle 13.5% and buffaloes 9.3%), respectively. The potential risk factors associated with brucellosis were studied. The iELISA seroprevalence in aborted, pregnant and lactating animals were 31.6%, 13.5% and 21.1%, respectively and in animals with repeated breeding, retained placenta and birth of weak calves were 11.3%, 12.3% and 15.3%, respectively. Our results provide baseline information on the epidemiology of bovine brucellosis and its potential risk factors in Punjab, Pakistan, which will ultimately help to improve control programs in the country. The higher seroprevalence of brucellosis indicates its relevance for public health.

**Keywords:** Bovine tuberculosis, seroprevalance, Milk, ELISA

## CRIMEAN CONGO HEMORRHAGIC FEVER (CCHF) IN THE SCENARIO OF DSP-BIORISK MANAGEMENT IN PAKISTAN

P.R. Nasrullah, M.A. Kakar\*<sup>1</sup>, Azizullah<sup>2</sup>, M.S. Qureshi<sup>3</sup> and Mithat Direk<sup>4</sup>

Department of Human Nutrition, University of Agriculture, Khyber Pakhtunkhwa, Pakistan; <sup>1</sup>The World Bank Pakistan, 18-B, Jinnah Town, Samungli Road, Quetta, Pakistan; <sup>2</sup>Livestock and Dairy Development Department Balochistan Quetta, Pakistan

<sup>3</sup>Faculty of Veterinary and Animal Sciences University of Agriculture, Khyber Pakhtunkhwa, Pakistan; <sup>4</sup>Department of Agricultural Economy, Faculty of Agriculture, Selçuk University, Konya, Turkey

[adaqul@yahoo.com](mailto:adaqul@yahoo.com)

### ABSTRACT

The present condition of Crimean-Congo hemorrhagic fever is deteriorating in Pakistan. The tick-borne Nairovirus causing Crimean Congo Hemorrhagic Fever (CCHF) has taken a greater death toll on human as well as cattle, compared to last year. To bring to light, the disease, first time activated during the 1944 in Crimea killing 200 army officers of the Soviet Union, has now a vast geographical dispersal becoming endemic in 30 countries across the globe which includes Africa, Middle East, Asia and the Balkans. Ticks from the Bunyaviridae family especially the genus *Hyalomma* known to be primarily involved in its spread and an active carrier of the virus, cause it among animals and humans by a single bite, while the infected animals through skin contact, slaughter houses, veterinarians and other agricultural workers, expose the incidence of the disease to other animals and humans. Similarly, human-to-human transmissions can occur by blood contact, exchange of bodily fluids and secretions of the infected person, less precautionary hospital environment with CCHF patients, and mishandling and re-use of needles or other medical appliances. Though, the symptoms are not immediately visible by human-to-human blood contact (5 to 6 days), the tick-bite within a day or few (1 to 7 days) causes fever, joint pain, headache, vomiting, dizziness, diarrhea, red spots on the upper mouth palate and sore throat. Treatment includes supportive care of symptoms and eliminating secondary infections. Even though, the drug Ribavirin is the lone FDA approved cure so far, yet, there is no vaccine made, to shield the spread of CCHFV, besides, the tick spell on animal followed by disease incidence, to the transmission to other ticks, usually goes undetected. The rudimentary cause behind the highest prevalence of CCHF in Balochistan is their not only their local cattle, but also, poor quarantine laws and health investigation on the borders with Iran and Afghanistan which are the top possible route of virus dispersal across border lines. 58% of CCHF positive patients of Balochistan have prospectively reported to have been to remote areas of Afghanistan, in parallel, 13 animals out of 21 randomly selected animals from Afghanistan and Iran's neighboring areas after ELISA tests revealed to be CCHF positive. Precautionary measures including medicinal care of cattle only last about a week or two which is worth than having no cure at all, on the other hand, human infections can be prevented by following the DSP Biorisk Management safety measures to avoid exposure to the disease and raising responsiveness on district and provincial level and acknowledge the population about this global concern before it becomes an epidemic.

**Key words:** CCHF, DSP, Bio-risk Management, Zoonotic Diseases, Pakistan.

## QUALITY ASSESMENT OF PASTEURIZED MILK THROUGH THE INCIDENCE OF SPORE FORMER BACTERIA IN PASTEURIZED MILK

Muhammad Asim Ikram, Imran Javed, Saima Inayat, Khalid Javed<sup>1</sup>, M. Junaid, H. Arslan Amin, Usman Mir Khan

Department of Dairy Technology, <sup>1</sup>Department of Livestock Production, University of Veterinary and Animal Sciences, Ravi Campus, Pattoki.

[asim.ikram@uvas.edu.pk](mailto:asim.ikram@uvas.edu.pk)

### ABSTRACT

Milk is a complete nutritious diet with very short shelf life. In order to extend the shelf life of milk Pasteurization is mainly used. Pasteurization is used to kill pathogenic organisms from milk and make it safer for human consumption. In this process, spore former bacteria (especially genus *Bacillus*, *Paenibacillus* and *Clostridium*) survive and they shorten the shelf life of milk. In order to access the microbial quality of pasteurized milk available in local market of Lahore, milk brands were analyzed for the enumeration of psychrotrophic, mesophilic and thermophilic spore formers (*Bacillus* and *Clostridium*) and total plate count. Total spore count was obtained by combining these and further identification was done by spore staining, morphological, biochemical identification tests. Fifty four samples of six available brands (A,B,C,D,E and F) were collected for the enumeration of spore former and total plate count. Spore former bacteria were cultivated using plate agar method on selective and mixed agar. The data was subjected to analysis through One Way Analysis of Variance (ANOVA) using DMR. Milk brand D showed less numbers of Total Plate Count (CFU/ml) as compared to other brands, while the thermophilic, mesophilic and psychrotrophic bacilli numbers were less in Brand E, A and D respectively. On the other hand the enumeration of thermophilic and mesophilic clostridia were less in Brand D in both cases. We can imagine the shelf life, extent of contamination and level of cross contamination of pasteurized milk, which is being sold in the local market of Lahore, with number of total spore former and Total plate count.

**Key Words:** Pasteurized milk, total plate count, spore former microbes



## CHALLENGES AND FUTURE SCOPE OF OSTRICH (*STRUTHIO CAMELIUS*) FARMING IN PAKISTAN

Muhammad Fiaz Qamar, Muhammad Arfan Zaman, and Kazim Ali

College of Veterinary & Animal Sciences, Sub campus: University College of Veterinary & Animal Sciences, Lahore.

[fiaz.qamar@uvas.edu.pk](mailto:fiaz.qamar@uvas.edu.pk)

### ABSTRACT

The ostrich (*Struthio camelius*) is a flightless bird and it has different names in various parts of the world, i.e, (Arabic na-ama, French autruche, German Straufi, Italian struzzo, Norwegian struts, Spanish avestruz). The ostrich has potential to provide human beings with low cholesterol and high quality food, utensils, clothing, and adornment; now a days there is a scope of ostrich farming for the procurement of meat, feathers, eggs and skin. There is a huge potential of extensive ostrich farming in Pakistan because abundant facilities regarding building design, bedding materials, fencing, environmental influences including sunlight are available throughout Urban and Rural areas of Pakistan. The breeding system needs special attention from academia and industry to study and interpret factors that influence hatchability. The major diseases and their management include Ostrich parasites, i.e, Ostrich wireworm (or stomach worm), Nematodes, tapeworm, protozoa, eye fluke and external parasites, Bacterial diseases; Botulism, Necrotic (or ulcerative) enteritis, Omphalitis and yolk sacculitis, Campylobacteriosis and Anthrax, Viral diseases; Avian influenza, Newcastle disease, Avian pox, Mycotic diseases: Candidiasis, Aspergillosis, Nutritional disorders: vitamin A, vitamin D, vitamin E, vitamin B, manganese, Economics/profits of ostrich farming comes through sale of ostrich eggs, sale of live ostriches and sale of ostrich products. Future of the ostrich industry in Pakistan is bright as there is a large scale market size, established leather processing and manufacturing, low feed cost per ostrich, best feed-to-weight ratio, low labour cost. Most Pakistanis like meat including farming based chicken, new and innovative investment opportunity for investors, potential for high production levels of breeder birds, potential for good food conversion efficiency in slaughter birds, Longevity of breeder birds leads to long-term commitment of producers to the future of the farmed ostrich.

Keywords: Ostrich Farming, Pros and Cons, diseases,



**ESTIMATION OF COMPOSITION, ADULTERATION AND MICROBIAL LOAD IN RAW MILK SAMPLES COLLECTED FROM DAIRY FARMS AND URBAN AREAS OF LAHORE DISTRICT, PAKISTAN**

Atta Muhammad Arif, Muhammad Ali<sup>1</sup>

University of Veterinary and animal sciences Lahore-Pakistan, <sup>1</sup>Department of Dairy Technology

[Muhammadali.uvas@gmail.com](mailto:Muhammadali.uvas@gmail.com)

**ABSTRACT**

Milk is a translucent white liquid rich in nutritional profile from mammary glands of animals which make it perfect source diet required for growth and maintenance in infants, children and adults. But for sake of higher profit it is severely polluted through poor processing and hazardous chemicals which ensure it a birthplace of many diseases. Three hundred and sixty raw milk samples were collected from urban areas (ten areas of Lahore city) and dairy farms (ten dairy farms located in Lahore) with an interval of two months. Samples were analyzed for the estimation of adulterants, chemical composition and total microbial load. The results indicated that added water was positive for 100% samples of urban areas and no sample was found positive for the added water for Dairy Farms. Adulterants like Formalin, Sugar and Urea were not detected in any sample while Starch, added water, Hydrogen peroxide, CO<sub>3</sub>-/HCO<sub>3</sub> and NaCl were detected in 4.667, 32.555, 2.00, 22.667, and 55.333% samples of raw milk from urban areas respectively. None of the adulterants were detected in dairy farm milk samples. The tests carried out for evaluating the chemical composition and physico-chemical properties of the raw milk samples from urban areas showed the following mean values of 2.758±0.288, 2.629±0.180, 1.270±0.077, 7.133±0.302, 6.456±0.264, 0.472±0.022 for lactose, fat, protein, TS, pH, ash respectively. Milk samples from the dairy farms recoded mean values of 4.816±0.321, 5.023±0.385, 3.363±0.180, 13.188±0.656, 6.669±0.265, 0.749±0.051 for lactose, fat, protein, TS, pH, ash respectively. Mean values of total plate count was 5.206±0.283 cfu/ml for urban areas raw milk samples and 4.674±0.273 cfu/ml for the dairy farm milk samples. The results depicted that raw milk samples analyzed from urban areas did not conform to even the minimum legal standards and possess adulterant chemicals which are noxious to health.

**Keywords:** Milk, adulteration, Lahore

## ANTHELMINTIC AND ANTIPROTOZOAL ACTIVITIES OF CALOTROPIS PROCERA UNDER FIELD CONDITIONS

Muhammad Ashraf, A. Javeed, M. Adil Rasheed, Qamar Niaz and Mukhtar Ahmad,  
Department of Pharmacology and Toxicology UVAS, Lahore  
[drashraf2001@uvas.edu.pk](mailto:drashraf2001@uvas.edu.pk)

### ABSTRACT

Plants are important sources of drugs and have been used for the treatment of various ailments in human and animals. Antipyretic, analgesic, anti-inflammatory, antispasmodic, antidiarrheal and anticoccidial effects of *Calotropis procera* leaves have been reported. Presences of cardiac glycosides, alkaloids, flavonoids and triterpenes have been detected in its different parts. Anthelmintic activity of *C. procera* leaves was evaluated against nematodes of range land sheep under field conditions. Parasitic burden was evaluated by counting eggs per gram (EPG) of faces by McMaster technique before and after medication. Moxidectin (0.2mg/kg I/M) was used as positive control. Chloroform extract of *C. procera* leaves was administered at various doses (3, 6 and 12mg/kg) orally. Dose depended effect in the reduction of EPG was observed 91.4 % at the dose of 12mg/kg at day 14 post medication while with Moxidectin it was 93.3 %. No significant differences were observed in the renal (creatinine, urea) and liver function (ALT, AST, ALP) tests which were in the normal range in groups of *C. procera* and moxidectin treated animals. Theileriosis is an important blood protozoal disease of ruminants and leads to heavy mortality. Infected animals were diagnosed and confirmed by PCR and treated with the chloroform extract of *Calotropis procera* leaves (3mg/kg IM ) or of *Peganum harmala* leaves (3mg/kg IM). Buparvaquine preparation (Butalex 2.5 mg/kg IM) was used as positive control drug. Anti-theilerial efficacy of treatments was estimated by evaluating the clinical manifestation of disease and parasitological findings. Beside this treatment effect on hematological and biochemical reactions of liver and kidney functions were determined. It was found that animals treated with *C. procera* chloroform extract and butalex had rectal temperature in normal range by the day 7 of post-treatment. A non-significant difference ( $P>0.05$ ) in schizonts and piroplasms in various treatments were observed at 14 and 21 day of post treatment.

**Keywords:** *C. procera*, chloroform extract, anthelmintic activity, antiprotozoal activity

**TOXICOLOGICAL EFFECTS OF AFLATOXIN B1 ON GROWTH PERFORMANCE, HUMORAL IMMUNE RESPONSE AND BLOOD PROFILE OF JAPANESE QUAIL**

Muhammad Younus, Sakhra Mahmood, Asim Aslam, Aftab Ahmed Anjum, Aman Ullah Khan, Asif Idrees, M.L. Sohail

University of Veterinary & Animal Sciences, Lahore-Pakistan

[younusrana@hotmail.com](mailto:younusrana@hotmail.com)

**ABSTRACT**

Toxic effects of Aflatoxin B1 on growth performance and humoral immune response against Newcastle disease Virus (NDV) vaccine were determined in Japanese quails. Quail chicks (n=300) were divided into five groups and each group was further subdivided into two subgroups. One subgroup of each group was vaccinated with NDV vaccine to study the antibody response. Experimental diets were prepared by adding different levels of AFB1 i.e. 0.25mg/kg feed, 0.5mg/kg feed, 1 mg/kg feed and 2 mg/kg feed, and were offered to groups B, C, D and E respectively for 42 days, while group A was kept as control. Feed intake, body weight gain and feed conversion ratio were calculated on weekly basis. The Antibody titers of vaccinated quails were measured by haemagglutination inhibition test. Hematological parameters Erythrocyte sedimentation rate, Hb content, hematocrit and total leukocyte count were measured in AFB1 fed birds and were compared with the quails given basal diet. Results showed significant reduction ( $P \leq 0.05$ ) in feed intake, weight gain and feed conversion ratio, when compared to control group. Moreover, anti-NDV antibody titre was significantly reduced ( $P \leq 0.05$ ) in AFB1 fed groups. Hb content, hematocrit, and TLC had time and dose dependent decrease in AFB1 fed quails while ESR was increased significantly in AFB1 fed birds. This highlighted adverse effects of AFB1 on growth performance, humoral immune response and blood profile of quails.

**Keywords:** Toxicology, aflatoxin, immunity

**DISPOSITION KINETICS OF CIPROFLOXACIN AFTER INTRAMUSCULAR ADMINISTRATION IN LOHI SHEEP**Zahid Iqbal<sup>1</sup>, Ijaz Javed<sup>2</sup>, Riaz Hussain<sup>3</sup>, Mudassar Mohiuddin<sup>3</sup>

1Department of Pharmacology, Isra University, Islamabad, 44000, Pakistan

2Institute Pharmacy, Physiology and Pharmacology, University of Agriculture, Faisalabad, 38040, Pakistan; 3Department of Pathobiology, College of Veterinary and Animal Sciences, The Islamia University, Bahawalpur, 63100, Pakistan

[zahid1.iqbal1@gmail.com](mailto:zahid1.iqbal1@gmail.com)**ABSTRACT**

Disposition kinetics of ciprofloxacin studied in domestic Lohi sheep. The purpose was to optimize dosage regimen of the drug in local animals. 5 mg/Kg ciprofloxacin intramuscular (IM) preparation was injected to the thigh muscle of eight animals of similar age and weight. In every animal, a blank blood sample was taken before injecting the drug. After start of experiment, blood samples were collected from each animal at various time periods for estimating the concentration of ciprofloxacin in the blood using an advanced HPLC technique. An open two compartmental pharmacokinetic model was applied to calculate different disposition parameters. In this study the absorption half life ( $t_{1/2 \text{ abs}}$ ) was found to be  $0.63 \pm 0.16$  hours while maximum concentration in plasma at any time,  $C_{\text{max}}$ , was observed as  $1.97 \pm 0.15$   $\mu\text{g/ml}$ . Time taken by drug to reach maximum concentration in the plasma ' $T_{\text{max}}$ ' was observed at  $0.88 \pm 0.09$  hours. Distribution and elimination half life values were observed to be  $0.46 \pm 0.05$  and  $2.93 \pm 0.45$  hours, respectively. A higher volume of distribution ( $V_d$ ) as  $2.89 \pm 0.30$  L/kg was seen in this study. Total body clearance (CL) was observed as  $0.75 \pm 0.04$  L/hr/kg and while area under the plasma concentration versus time curve (AUC) showed value of  $7.19 \pm 0.38$   $\mu\text{g}\cdot\text{hr/mL}$ . These calculated disposition parameters were further used to optimize IM dosage of ciprofloxacin in local sheep. We concluded that in local sheep, a higher dosage (21.43 mg/kg) should be used intramuscularly and be repeated after 24 hours interval so that antimicrobial resistance may be avoided.

**Keywords:** Lohi sheep, Ciprofloxacin, Disposition kinetics, Optimal dosage

**EFFECTS OF DIETARY VITAMIN C ON ARSENIC INDUCED TOXICITY ON SEMEN QUALITY, REPRODUCTIVE HORMONES AND HISTOPATHOLOGY OF TESTES IN TEDDY GOAT BUCKS**

<sup>1</sup>Muhammad Zubair, <sup>2</sup>Maqbool Ahmad, <sup>3</sup>Mushtaq Ahmad, <sup>2</sup>Saqib Umar, <sup>2</sup>Shujait Ali

<sup>1</sup>Faculty of Veterinary Sciences University of Poonch Rawalakot Azad Kashmir.

<sup>3</sup>University of Veterinary and Animal Sciences Lahore; <sup>2</sup>Department of Theriogenology, University of Agriculture Faisalabad, Pakistan

[drzubairabbasi@gmail.com](mailto:drzubairabbasi@gmail.com)

**ABSTRACT**

Arsenic is considered as one of the major environmental toxicant affecting male reproductive system. The present study was conducted to investigate the toxic effects of arsenic on semen quality and reproductive hormones of Teddy bucks. Vitamin C was also fed to find the ameliorative effects on these parameters. For this purpose, twelve adult Teddy bucks were randomly divided into three treatment groups: A (control), B (Arsenic 5 mg/kg BW/day orally) and C (Arsenic as in group B and vitamin C with oral dose of 200 mg/kg BW/day). These treatments were continued for 12 weeks. Semen quality parameters (volume, motility, count, sperm morphology, sperm membrane integrity, sperm DNA integrity and acrosomal integrity) of experimental bucks of each group were evaluated on weekly basis. The libido of each buck was also observed before the semen collection. Blood was collected after every two weeks to measure the serum male hormones (testosterone, leutinizing hormone and follicle stimulating hormone). Malondialdehyde and cortisol were also evaluated after every two weeks. At the end of the study, testes of all the animals were collected and evaluated for histopathological studies. The obtained data was subjected to two-way analysis of variance, followed by Tukey test for multiple mean comparisons. The intensity of libido and semen evaluation parameters in arsenic treated buck were significantly ( $p < 0.05$ ) reduced. The level of male hormones (testosterone, leutinizing hormone and follicle stimulating hormone) were also significantly ( $p < 0.05$ ) decreased in arsenic treated animals. However, the values of malondialdehyde and cortisol were increased significantly in toxic group. The histopathological lesions in testes were present in the form of loss of germinal epithelium and atrophy of leydig cells. It was concluded that the supplementation of vitamin C ameliorated the toxic effects of arsenic on semen quality, hormones, malondialdehyde and cortisol.

**Keywords:** vitamin C, arsenic, toxicology, hormone

## CHARACTERISTICS OF MEAT EMULSION SYSTEMS WITH BEEF REPLACED BY SOY PRODUCTS

Neslihan Polat, Mustafa Karakaya, Ali Samet Babaoğlu, Kübra Ünal.

Agriculture Faculty, Food Engineering Department, Selcuk University, Konya, Turkey  
[asmtbb@gmail.com](mailto:asmtbb@gmail.com)

### ABSTRACT

Soy products can be used in the meat products to improve emulsion and textural properties, enhance nutritional value and reduce cost of production. The aim of this study was to evaluate the effect of addition of soy protein products at four concentrations on the technological properties (water holding capacity and cooking loss). Minced meats were divided into twelve treatment groups. Three soy protein products (tofu, textured soy protein (TSP), soy protein isolate (SPI)) and four concentrations (0%, 25%, 50% and 75%) of these compounds were added to meat samples, and homogenized, respectively. Samples in control treatment (0%) had no soy protein products added. It was also determined that the effect of soy protein products and their concentrations on pH and emulsion properties (emulsion capacity, emulsion stability and color parameters) of O/W emulsions prepared from beef meat. The highest water holding capacity and the lowest cooking loss were reached when 75% of soy products was added. Addition of textured soy protein increased water holding capacity and decreased cooking loss of beef meat samples. However increase in the concentration of soy products decreased emulsion capacity, emulsion stability, L\* and a\* values of emulsions.

**Key words:** soy products, beef meat, emulsion properties

**AFLATOXIN M1 PREVALENCE IN MILK AND RECENT UPDATES ON DECONTAMINATION STRATEGIES**

Dr. Amir Ismail, Dr. Muhammad Riaz, Dr. Saeed Akhtar

Institute of Food Science and Nutrition, Bahauddin Zakariya University, Multan, Pakistan

[amirismail@bzu.edu.pk](mailto:amirismail@bzu.edu.pk)

**ABSTRACT**

Aflatoxin M<sub>1</sub> (AFM<sub>1</sub>) is a group 2B category carcinogen according to International Agency for Research on Cancer and is of most concern for humans because of its prevalence in milk and milk products. Dairy products are of most concern for health authorities because of the fact that these food commodities are consumed maximally by the infants and elderly, the two age groups having minimum levels of immunity. Global data indicates the prevalence of AFM<sub>1</sub> above permissible limits especially in the developing countries suggesting the need for concerted efforts to curtail its level in dairy products. The best approach to control this toxin is to prevent dairy feed from aflatoxin producing fungus, however the curative approaches are also under research to reduce the level of AFM<sub>1</sub> below cut off values for human health. More recently, microorganism especially lactic acid bacteria are employed for binding of AFM<sub>1</sub> in order to reduce the bioavailability of AFM<sub>1</sub>. The use of clay materials is also reported to reduce AFM<sub>1</sub> levels without disturbing the general characteristics of milk. Present review will provide recent data regarding the prevalence of AFM<sub>1</sub> in milk and milk products, their health impacts, regulations adopted in different countries, methods reported to have potential to reduce the level of AFM<sub>1</sub> in milk and the complications involved in the use of these reduction methods.

**Keywords:** Aflatoxin, milk, health

## COLOSTRUM WHEY: A POTENTIAL INGREDIENT FOR DAIRY SECTOR IN NUTRITIONAL ASPECTS

Maryam Batool, Saima Inayat, Muhammad Asim Ikram

University of Veterinary and Animal Sciences Lahore, Pakistan  
[maryambatool17@gmail.com](mailto:maryambatool17@gmail.com)

### Abstract

Whey protein is a well-know nutritional supplement in the sports arena, including body building. Colostrum whey is the liquid remaining after the removal of casein and fat. Colostrum whey is very nutritional as it is probiotic naturally. Bovine Colostrum whey in particular contains more than 200 different proteins, with  $\alpha$ -lactalbumin,  $\beta$ -lactoglobulin, bovine serum albumin (BSA), and immunoglobulins as the major constituents. Bulk bovine whey has been used in the production of nutraceuticals. Colostrum whey contains compounds with a direct bactericidal effect. In addition, it also contains compounds that trigger immune defense mechanism to further help in elimination of bacteria and viruses. It will be beneficial if colostrum whey is utilized to convert into naturally nutritional by products such as ricotta cheese etc. Colostrum whey average values of fat, protein, TS, ash and chlorides are  $0.333 + 0.133$ ,  $2.850+0.029$ ,  $7.257 + 0.139$ ,  $0.070 + 0.012$  and  $0.320 + 0.006$ . It contains different bioactive components which will be beneficial in a result of neutraceutical byproducts with natural probiotic nature which will help to improve our internal microflora for better digestibility and in increasing immunity. Now a days, Wastage of whey is a big problem for the environment due to high BOD value. So it would be beneficial, if colostrum whey is converted into value added product as it is a natural gift of God. Conversion of colostrum whey into cheese or energy drinks would give an innovative route to the Dairy Industry future.

**Keyword:** Colostrum whey, Colostrum whey benefits, Colostrum whey products



**DIFFERENTIAL GROWTH OF SALMONELLA PULLORUM AND SALMONELLA GALLINARUM USING MODIFIED SS AGAR AND RV MEDIUM**

Bakhtawar Maqbool and Sajjad-ur-Rahman  
University of Agriculture, Faisalabad, Pakistan  
[bakhtawardogar@yahoo.com](mailto:bakhtawardogar@yahoo.com)

**Abstract**

Salmonella belongs to family Enterobacteriaceae and most of the serovars are considered to be zoonotic and food borne pathogenesis. Salmonellosis is generally diagnosed through the culture techniques which take more time (4-7days). Many rapid diagnostic methods are also available in the market like latex agglutination test, enzymes linked immunosorbant assay( ELISA) and polymerase chain reaction ( PCR) which are used but the problem is that they are lacking specificity. So, the present study was planned to differentiate S. pullorum and S. gallinarum by using modified Salmonella Shigella (SS) agar and Rapport Vassilidas medium. Thirty five samples were collected from the intestinal content of poultry birds and were diluted with normal saline, further enrichment was performed using RV broth. Furthermore, subculturing was done on Macconkey and SS agar. The biochemical tests were performed using Rapid ID ONE kit, this kit contains all the biochemical tests necessary for species level identification. Polymerase chain reaction was employed for strain level identification, in which the rat A gene was targeted. Out of the 35 samples 1(2.85%) positive for S. pullorum 5.71% for S. Gallinarum. Various concentrations of dulcitol (D) were used in RV medium and SS agar. In RV medium the dulcitol concentration was 0.5g (D), 1g (D) and 0.5 g (D) + 0.5g lactose. For SS agar 0.5g (D), 1g (D) and 0.5 g (D) + 0.5 g lactose. Results show that with 1g concentration of dulcitol shows better growth in both RV and SS agar for S. gallinarum. While in 0.5 g (D) + 0.5 g lactose the growth of S. Pullorum was totally inhibited. This research provides an insight to how the routinely used growth medias for Salmonella species may be modified with regards to specific ingredients, and further modifications may be made by microbiologists for improved culturing results.

**Key words:** Modified medium, SS Agar, Salmonella, differential diagnosis

## COMPARATIVE RESILIENCE OF TEDDY AND BEETAL GOAT BREEDS AGAINST ARTIFICIAL INFECTION OF HAEMONCHUS CONTORTUS THROUGH IMMUNOGLOBULINS AND HISTAMINE

Asim Shamim<sup>1\*</sup>, Muhammad Sohail Sajid<sup>2,3</sup>, Muhammad Nisar Khan<sup>2</sup>, Muhammad Saqib<sup>4</sup>, Muhammad Arfan Zaman<sup>5</sup>, Muhammad Imran<sup>2</sup>, Hafiz Muhammad Rizwan<sup>2</sup>

<sup>1</sup>Department of Pathobiology, University of Poonch Rawalakot Azad Kashmir, Pakistan;

<sup>2,3</sup>U.S. Pakistan Center for Advanced Studies in Agriculture and Food Security (USPCAS-AFS); <sup>4</sup>Department of Veterinary Clinical Medicine and Surgery, University of Agriculture, Faisalabad, Punjab, Pakistan; <sup>5</sup>Department of Pathobiology, College of Veterinary and Animal Sciences, University of Veterinary and Animal Science Lahore.

[asimshamimuajk@gmail.com](mailto:asimshamimuajk@gmail.com)

### Abstract

Haemonchus (H.) contortus, an abomasal worm of goats (*Capra hircus*), affects health, production and welfare globally. Usually controlled through anthelmintics; however, development of anthelmintic resistance (AR) and concern of drug residues in food products makes use of anthelmintics questionable. Evaluation and selection of goat breeds towards Haemonchus infection is an alternate approach to control this parasite. Objective of this study was to determine the comparative resistance / resilience of Teddy and Beetal goat breeds of Pakistan of immunoglobulin (Ig) isotypes and histamine concentrations in artificially infected goats with third stage larvae (L3) of H. contortus. Experimental goats (n=60) were divided into two groups i.e. Teddy and Beetal each divided into six sub-groups containing 5 goats in each sub-group. Two infection protocols were followed to infect goats i.e. early and late. Goats received dose of 12000 and 18000 L3 of H. contortus. Adult H. contortus antigen was prepared for performing enzyme linked immunosorbent assay (ELISA). Indirect ELISA was performed to evaluate the response of Igs and histamine against H. contortus. There was significant (P<0.05) difference observed in the levels of IgG and IgE at different time intervals post infection. Concentration of histamine was persistent in the plasma of experimentally infected goat breeds throughout the trial. However, it was significantly different at 4th and 6th weeks post infection. Up and down regulation of Igs and histamine post infection makes Teddy goats better responder than Beetal goats. Further investigations on mechanisms involved in breed resilience and/or susceptibility are required.

**Keywords:** Haemonchus contortus, Teddy, Beetal, Goats, Histamine, immunoglobulins, ELISA.

**HIGH PROTEIN DIET INDUCES B-CELL REGENERATION AND RESTORES NORMAL GLYCAEMIA IN ALLOXAN INDUCED DIABETIC RATS**

Humaira Muzaffar\*, Muhammad Naeem Faisal, Tanweer Khaliq, Junaid Ali Khan, Bilal Aslam, Aisha Mahmood

Institute of Pharmacy, Physiology and Pharmacology, University of Agriculture Faisalabad, Pakistan

[2007ag1556@uaf.edu.pk](mailto:2007ag1556@uaf.edu.pk)

**Abstract**

Diet is a key factor in promoting health, metabolic homeostasis and fulfilling energy requirements. Diabetes mellitus is a metabolic disorder and characterized by hyperglycaemia resulting from defect in insulin production or secretion, insulin action or both. So,  $\beta$ -Cell dysfunction is a critical component in the development of diabetes. The amount, type and balance of the main dietary macronutrients (carbohydrates, proteins and fats) have a great impact on the management of diabetes. However, little is known about the molecular mechanism that how high protein diet is responsible for functional changes in pancreatic  $\beta$ -cells. This study was designed to investigate the effect of high protein diet on physical parameters, glycaemic control,  $\beta$ -cell regeneration, insulin production and release in diabetic rats. High protein diet was administered to alloxan induced diabetic rat model. Blood samples were taken for measuring blood glucose level, lipid profile and other biochemical tests. Tissue samples of pancreas and liver were collected for morphological study. Results indicated that high protein diet counteracted the reactive oxygen species (ROS) produced inside  $\beta$ -cells and normalized the elevated serum glucose, triglycerides and cholesterol levels and helps in normalizing the pancreatic tissue structure, morphology as well as normal replenishing or regeneration of  $\beta$ -cells. The data was analyzed statistically through one way ANOVA and DMR.

**Keywords:** high protein diet, alloxan induced diabetes, glycaemic control,  $\beta$  cell regeneration.

**ANTIMICROBIAL POTENTIAL OF BOVINE LACTOFERRIN AGAINST  
FOODBORNE PATHOGENS**

Ishtiaque Ahmad, Ammarah Khatoun, Muhammad Ayaz, Muhammad Asim Ikram,  
Muhammad Junaid, Usman Mir Khan, Nisar Ahmad

University of Veterinary and Animal Sciences, Lahore, Pakistan

[ishtiaque@uvas.edu.pk](mailto:ishtiaque@uvas.edu.pk)

**Abstract**

Food born illnesses are always a serious health issue in the Pakistan and throughout the world. Lactoferrin can be isolated from cheese whey by cation exchange chromatography without loss of its biological properties. Lactoferrin exhibits strong antimicrobial activity against different bacteria, virus, protozoa, fungi and yeast. The present study is concerned with the antibacterial activity and sensitivity of bovine lactoferrin against foodborne pathogens and to evaluate the pH range and thermal stability of bovine lactoferrin. Agar gel diffusion test was used to check antibacterial activity for each pathogen. Bacterial samples were purified and confirmed by bio-chemical tests. Three concentrations of lactoferrin 0.5mg/ml, 1mg/ml and 10mg/ml were treated on Escherichia coli, Salmonella enteritidis and Staphylococcus aureus. Antimicrobial activity of lactoferrin was determined at different range of pH. The heat stability of lactoferrin was determined at different heat treatments. The data was analyzed by complete randomized design (1-factor factorial) and comparison of means was done by Duncan's multiple range test. Outcomes received indicated that all three pathogenic bacteria were inhibited, even at 0.5mg/ml significantly. Escherichia coli is most sensitive and Staphylococcus aureus is least. Lactoferrin remain effective at 6, 7, 8 and 9 pH but no inhibition showed at 3, 4 and 5 pH significantly. There is no significant effect on antibacterial activity at pasteurization temperature i.e. 60-75°C but destroyed above 80°C. The results suggested that supplementation of lactoferrin in different food products enhances the food safety and quality.

**Key words:** lactoferrin, bovine, foodborne pathogens, antimicrobial activity

**THE EFFECTS OF KEFIR IN TENDERIZING CHICKEN AND TURKEY BREAST MEAT**

Kübra Ünal, Ali Samet Babaoğlu, Mustafa Karakaya  
Agriculture Faculty, Selçuk University, Konya, Turkey  
[ulusoy\\_kubra@hotmail.com](mailto:ulusoy_kubra@hotmail.com)

**Abstract**

Tenderness is the most quality properties of consumer acceptability of meat and meat products. The aim of this study was to evaluate the usage of kefir on some tenderness properties, pH and color characteristics of chicken and turkey breast meats. Meat samples were marinated in kefir for 24, 48 and 72 hours. Instrumental tenderness of cooked breast meat samples was measured using the Warner Braztler Shear blade. Marinating with kefir was decreased lightness of samples.  $a^*$  values of samples increased with using kefir.  $L^*$  and  $a^*$  values of marinated chicken and turkey breast meats were, 60.43, 65.18; 11.33, 8.96. There was no difference ( $p < 0.05$ ) in  $b^*$  values between marinated and non-marinated samples. The pH of marinated samples was decreased according to the control group. pH values of marinated chicken and turkey breast meat were 5.52 and 5.49, respectively. Firmness of turkey breast meat decreased with usage of kefir, while marinated with kefir did not change the firmness of chicken breast meat.

**Key words:** kefir, tenderness, chicken meat, turkey meat.

## COMPARATIVE EFFICACY OF SYNCHRONIZATION PROTOCOLS (PROGESTERONE BASED OR FREE) ON CONCEPTION RATE IN CAPRINE SPECIES

Maqbool Ahmad, Umair Asharaf, Shafia Tehseen Gul, Rehana Kousar, Ijaz Ahmad, Khurram Ashfaq

University of Agriculture, Faisalabad, Pakistan

[drmaqbool\\_ahmad@yahoo.com](mailto:drmaqbool_ahmad@yahoo.com)

### Abstract

This study was designed to investigate the effect of different exogenous hormonal strategies for improving the fertility in goats. The major objective was to evaluate the comparative efficacy of progesterone-based and progesterone-free estrus synchronization protocols on conception rate in dwarf goats. North Carolina Synchronization (NCSynch) is a new strategy for synchronization of estrous. A total of 27 does were divided into 3 equal groups A, B and C. Group-A was administered with slightly modified NCSynch protocol i.e., PGF2 $\alpha$  which was given on day 1 and 15, GnRH on day 8 & 17 of the experiment. Group-B was given medroxy-progesterone in combination with PGF2 $\alpha$  and GnRH. Sponges impregnated with medroxy-progesterone acetate were placed intra-vaginally for period of 15 days. Prostaglandin F2 $\alpha$  were administered upon sponge removal on day 15 of experiment and GnRH was given on day 17 to these animals. Group-C comprised of control animals receiving normal saline I/M on day 1, 7, 15 and 17. Both treatments were evaluated against control Group-C. The onset of standing estrus and duration of heat period were also recorded. Controlled natural mating was performed in all groups. Progesterone levels were also assessed during the trial by using radio immunoassay (RIA) on day 1, 7, 15 and 17 of experiment. Overall response to estrus was 55, 100 and 33% in Group-A, B and C respectively. Conception rate was 55, 88 and 33% in group A, B and C respectively. Mean level of progesterone was significantly different in pregnant animals of Group-A from than that of Group-B and C during treatment on days 1st, 7th & 15th.

**Key words:** Caprine, estrus synchronization, methods, progesterone, NCSynch

## WATER PRODUCTIVITY UNDER GOVERNMENT AND FARMERS CONTROLLED CANAL IRRIGATION SYSTEMS

Muhammad Jamal Khan<sup>1</sup>, Rabnawaz<sup>1</sup> and Amanullah<sup>2</sup>

Faculty of Crop Production Sciences, The University of Agriculture Peshawar, Pakistan-25130

[mjkhan@aup.edu.pk](mailto:mjkhan@aup.edu.pk)

### Abstract

Pakistan is becoming a water scarce country with declining per capita water availability (<1000 m<sup>3</sup>), whereas its demand for domestic, industrial and environment is rapidly increasing resulting in to stress on sustainable water supply to irrigated land contributing >90% of the agricultural production. Increase in crop water productivity is one of the most important options to achieve efficient and effective use of water resources in arid to semi-arid environment. The present study was conducted to compare the crop water productivity under public (Kabul River Canal) and civil (Joe Sheikh Canal) systems in Peshawar. Water productivity was determined at system as well as farm level for both the systems. Performance of two systems was assessed using twenty-three indicators in five major groups (productivity, water supply, sustainability, environment and institutional/management). During winter season (Rabi), wheat, sugarcane, vegetables, fodder, and orchards occupied 43, 24, 14, 6, 7% and 34, 31, 25, 6, 3% of the cropped area under public and civil canal systems, respectively. In summer season (Kharif), maize, sugarcane, vegetables, fodder, and orchards occupied 41, 24, 12, 10, 7% and 35, 31, 23, 5, 3%, land under public and civil canal systems, respectively. The cropping intensities of 187% and 194% were recorded in the two systems, respectively. Average crop water productivity (CWP) of wheat, maize, sugarcane and tomato were 0.96, 1.11, 3.31, 3.61 kg m<sup>-3</sup> under public and 0.90, 0.77, 2.38, 2.98 kg m<sup>-3</sup>, respectively. Despite huge amount of expenditure on management as well as operation and maintenance of public system, benefit-cost ratio of civil system was 40% high. The present study concludes that the civil system with minimal management and operation and maintenance cost performed better than public system. However, to further improve the productivity as well as performance, management of both the system needs to be gradually transferred to the farmers.

**Keywords:** Water conservation, crops, management, farmers

## SYNTHETIC POLYMORPHIC PEPTIDES: A STEP FORWARD FOR DIAGNOSIS OF TYPE-SPECIFIC TOXOPLASMOSIS IN HUMANS AND ANIMALS

H.M. Rizwan<sup>\*1,2</sup>, D. Arranz-Solís<sup>2</sup>, MS Sajid<sup>1,3</sup>, J.P. Saeij<sup>2</sup>

<sup>1</sup>Department of Parasitology, University of Agriculture, Faisalabad, Pakistan;

<sup>2</sup>Department of Pathology, Microbiology and Immunology, University of California, Davis, USA; <sup>3</sup>U.S.-Pakistan Center for Advanced Studies in Agriculture and Food Security, University of Agriculture, Faisalabad, Pakistan.

[dr.hmrizwan@yahoo.com](mailto:dr.hmrizwan@yahoo.com)

### Abstract

Toxoplasmosis, an important cosmopolitan zoonotic disease of humans and animals, is caused by an obligate intracellular parasite *Toxoplasma (T.) gondii*. Ingestion of uncooked or undercooked meat infected with viable tissue cysts and other foods infected with oocyst are major sources of disease transmission. *Toxoplasma gondii* has a clonal population structure containing three dominant clonal lineages, i.e. the clonal types I, II and III. However, some recombinant forms or exotic types also exist which develop from the combination of these clonal types. In previous studies, typing of *T. gondii* has been done through DNA based techniques. For this, DNA samples of *T. gondii* were collected either from oocysts collected from feline faecal samples or from tissues/tissue cysts collected from euthanized cats. However, getting an adequate amount of parasitic DNA from subclinical and clinical cases of toxoplasmosis remains bottleneck of DNA based typing techniques. Use of synthetic polymorphic peptides derived from *T. gondii* antigens is a simple, cost effective, sensitive and specific typing method. The results of previous studies showed that synthetic polymorphic peptides derived from ROP1, SAG1, GRA3, GRA6, GRA7 antigens discriminate type II from non-type II infections which suggests that serotyping is one of the promising methods for typing strains; although, limitations exist as a consequence of higher peptide polymorphism. Other peptides from different markers must be studied in order to discriminate clonal as well as non-clonal type.

**Key words:** Toxoplasmosis, Serotyping, Peptides, *Toxoplasma gondii*,



**FASCIOSIS: A MAJOR THREAD OF FOOD SECURITY AND ECONOMIC LOSSES ALL AROUND THE WORLD**

Hafiz Muhammad Rizwan<sup>\*1,2</sup>, Muhammad Sohail Sajid<sup>1,3</sup>, Muhammad Saqib<sup>4</sup>, Zafar Iqbal<sup>1</sup>

<sup>1</sup>Department of Parasitology, University of Agriculture, Faisalabad, Pakistan, <sup>2</sup>Department of Pathology, Microbiology and Immunology, University of California, Davis, USA. <sup>3</sup>U.S.-Pakistan Center for Advanced Studies in Agriculture and Food Security, <sup>4</sup>Department of Clinical Medicine & Surgery, University of Agriculture, Faisalabad, Pakistan.

[dr.hmrizwan@yahoo.com](mailto:dr.hmrizwan@yahoo.com)

**Abstract**

Of the parasitic diseases, helminth infections are the most common and the most economically significant diseases affecting grazing ruminants worldwide. Among these, fasciolosis, caused by *Fasciola* (*F.*) *hepatica* and *F. gigantica*, is one of the most significant and debilitating liver damaging diseases of ruminants. It has been estimated that about 67 hundred million ruminants are at risk of fasciolosis around the world. This disease can cause blood loss, reduced appetite, nitrogen retention, and impaired energy metabolism due to liver damage, reduction in weight gain and reduced milk production. Damage of organs and other biochemical changes are due to the migratory and hematophagic activities of *Fasciola*. The annual economic loss due to fascioliasis in the veterinary field was estimated US\$2–3 billion, as a result of liver condemnation, animal weight loss, reduction of milk yield and fertility declines. In Ethiopian Highlands the estimated economic losses due to fasciolosis were US\$2.75 million per year. The estimated losses due to this disease was reported US\$28 million in United Kingdom, in Bangladesh approximately US\$3 million in and in Pakistan US\$0.35 per year in terms of animal weight loss, declined fertility rate, and reduced milk yield. It is very difficult to control the disease and eradication is impossible as the parasite can survive in the host, snail, and environment for long periods. In addition to good farm management, it is imperative to exercise strategic anthelmintic therapy in order to maintain parasitic diseases at an acceptable level.

**Keywords:** Fasciolosis, Economic significance, Food security, Ruminants

**EFFECT OF BREED ON CARCASS TRAITS OF KUNDHI AND NILI RAVI BUFFALO**

Rameez Raja Kaleri, Hubdar Ali Kaleri, Muhammad Siddique Zardari  
Department of Animal Breeding & Genetics, Sindh Agriculture University, Tandojam  
[rameezkaleri@gmail.com](mailto:rameezkaleri@gmail.com)

**Abstract**

Present study was conducted to study the carcass traits of Kundhi and Nili Ravi buffalo. The total 100 animals of Kundhi and Nili Ravi breed were randomly selected and divided into A, B, C and D group. In group A and C there were Kundhi and Nili Ravi male whereas, B and D females of both breeds respectively and the age of animals was around 36 months. The selected animals were slaughtered at slaughter house of Seven Star meat processing company Dhabeji, Thatta. The carcass traits studied including live body weight, carcass weight, dressing percentage and boneless weight. The results revealed that carcass traits live body weight, carcass weight, dressing percentage and boneless weight of Nili Ravi male was significantly higher ( $P>0.05$ ) as compared to the Kundhi buffalo male, while same carcass traits were also observed highly significant ( $P>0.01$ ) in Nili Ravi female than Kundhi buffalo female. It was concluded that carcass traits of Nili Ravi buffalo are better expressed and produces more beef than Kundhi buffalo. Likewise Kundhi buffalo male is better in beef production than the Nili Ravi female and Kundhi buffalo female produces low carcass yield.

**Keywords:** Carcass traits, Breed, Kundhi, Nili Ravi buffalo.

**DAIRY SCIENCE PARK–EMPLOYMENT OPPORTUNITIES FOR THE YOUTH**

Sania Subhan Qureshi, Irfan ul Haq Qureshi, Sarzamin Khan and M Subhan Qureshi

Sania Subhan Qureshi

The University of Agriculture, Peshawar, Pakistan

[saniasq@aup.edu.pk](mailto:saniasq@aup.edu.pk)

**Abstract**

The Dairy Science Park (DSP) has emerged at the University of Agriculture, Peshawar, Pakistan, as a platform engaging the youth in productive economic activities. Medium-sized units (SMEs) are being networked with services providers, markets, and emerging entrepreneurs. The United Nations has accepted DSP as #SDGAction9671. Through the reshaped postgraduate research at the University, several entrepreneurship models were developed, supported with biorisk management. Several studies were conducted on feed conversion ratio, supplementation of antioxidants, growth pattern, breeding efficiency and effects of aflatoxins on economic parameters and other aspects. A study on effect of organic acids on the performance of Japanese quails found that net return was significantly higher by the supplementation. Artificial insemination in Japanese quails overcame fertility related problem as compared to natural mating. Proctodeal gland foam extender was found to be very effective for fertility, hatchability, sperm motility and count and as well as economical in terms of cost per chick. Analysis of the data generated on quails showed that a rolling fund of Rs.400,000 generated Rs.42,000 per month. Studies on rabbit farming system concluded that scope of rabbit as meat animal exists in the province and improvement in rabbit management could efficiently improve its production and utilization. Based upon the data generated so far, a feasibility of the production system has been prepared and we recommend it as an entrepreneur for the youth of the province. The data analysis revealed that an investment of Rs.200,000 has resulted in generation of a monthly income of Rs.41,325 per month through keeping 500 rabbit. The model was extended to the field through farmers training, especially sponsored by SRSP. The entrepreneurship models developed so far may be used for generating self employment for the youth and hygienic Halal food production for local consumption and export.

**Keywords:** Entrepreneurship, SMEs, SDGs, youth, livestock, poultry, value addition, quality control, biorisk management, export

## EFFECT OF RIPENING CONDITIONS ON PHYSICO-CHEMICAL PROPERTIES AND SURVIVAL OF PROBIOTIC MICROORGANISMS IN PROBIOTIC CHEDDAR CHEESE

Usman Mir Khan<sup>\*1</sup>, Fizzah Ashfaq<sup>3</sup>, Arslan Amin Bhatti<sup>1</sup>, Ishtiaque Ahmad<sup>1</sup>, Saima Inayat<sup>1</sup>, Muhammad Ayaz<sup>1</sup>, Nisar Ahmad<sup>2</sup>

<sup>1</sup>Department of Dairy Technology, University of Veterinary and Animal Sciences, Lahore, Pakistan; <sup>2</sup>Department of Livestock Production, University of Veterinary and Animal Sciences, Lahore, Pakistan; <sup>3</sup>Institute of Food Science and Nutrition, Bahauddin Zakariya Univesity, Multan, Pakistan  
[usmanmirkhan@yahoo.com](mailto:usmanmirkhan@yahoo.com)

### Abstract

Consumption of high concentration of probiotic bacteria at  $10^7$  viable cells per gram or per milli-litre of product is required to confer health benefits. Most publications concerning incorporation of probiotic bacteria into cheeses have focused only on their survival during manufacture and storage and only a few studies have considered the effect of probiotic adjuncts and the ripening temperatures on cheeses. Effect of ripening temperatures (4 and 8°C) on survival of probiotics microorganisms after ripening for 24 weeks. *Bifidobacterium animalis* subsp. *Lactis*, *Lactobacillus casei*, *Lactobacillus acidophilus* were used as an adjunct in the production of Cheddar cheeses which were ripened for 24 weeks at 4 and 8°C. The counts of starter lactococci in cheeses produced with *B. animalis*, *Lb. casei* or *Lb. acidophilus* ripened at 8 °C were significantly lower than those ripened at 4 °C at 24 weeks. Probiotic microorganisms remained viable ( $7.50 \log_{10}$  CFU/g) at the end of 24 weeks and their viability was not affected by the ripening temperatures. The acetic acid concentration in cheeses made with *Bifidobacterium* sp. or *Lb. casei* sp. was significantly higher than that of the control cheese. Increasing the ripening temperature from 4 °C to 8 °C did not affect the viability of the probiotic microorganisms, the salt, fat and protein contents of the cheeses during ripening of 24 weeks. This study shows that careful selection of probiotic microorganisms is required to have Cheddar cheese with high nutritional value for health benefits.

**Keywords:** Probiotics, Cheddar cheese, Ripening, Temperature, Nutritional Value.

**PHYSICO-CHEMICAL AND NUTRITIONAL PROPERTIES OF CHEDDAR CHEESE MADE FROM CITRUS RETICULATA BLANCO CRUDE FLOWERS EXTRACT (CFE)**

Usman Mir Khan<sup>\*1</sup>, Ishtiaque Ahmad<sup>1</sup>, Saima Inayat<sup>1</sup>, Muhammad Ayaz<sup>1</sup>, Nisar Ahmad<sup>2</sup>, Arslan Amin Bhatti<sup>1</sup>

<sup>1</sup>Department of Dairy Technology, University of Veterinary and Animal Sciences, Lahore, Pakistan; <sup>2</sup>Department of Livestock Production, University of Veterinary and Animal Sciences, Lahore, Pakistan

[usmanmirkhan@yahoo.com](mailto:usmanmirkhan@yahoo.com)

**Abstract**

*Citrus reticulata* Blanco crude flowers extract (CFE) at different level (1, 2, 3 and 4%) were used as natural milk coagulant instead of rennet to develop the Cheddar cheese. The Cheddar cheeses were compared with one controlled cheese made with rennet of 0.002%. Physico-chemical results of Cheddar cheese showed that cheese made with 1% and 2% of CFE had a longer and slightly softer texture. While, cheeses containing 3 and 4% CFE had semi-hard textural properties of curd similar to rennet added cheese. The CFE made cheese had moisture 37 %, fat 45 % on dry basis resembling to rennet made Cheddar cheese. Protein analysis shows that CFE made cheese had significant higher protein content than control one. The Cheddar cheese with 3% and 1% CFE were preferred by consumers instead of 2% and 4% for their better taste, texture/appearance and overall acceptability. Conclusively, it fulfills the nutritional requirement with acceptable organoleptic characteristics and at the same time provides health benefits.

**Keywords:** Cheddar cheese, *Citrus reticulata* Blanco, Buffalo milk, Milk coagulant.

**ENHANCEMENT OF OXIDATIVE STABILITY OF IRON ENRICHED BUTTER OIL WITH NATURAL ANTIOXIDANTS**Ikam Ullah<sup>1</sup>, Ishtiaque Ahmad<sup>1</sup>, Usman Mir Khan<sup>\*1</sup>, Nisar Ahmad<sup>2</sup><sup>1</sup>Department of Dairy Technology, University of Veterinary and Animal Sciences, Lahore, Pakistan; <sup>2</sup>Department of Livestock Production, University of Veterinary and Animal Sciences, Lahore, Pakistan[usmanmirkhan@yahoo.com](mailto:usmanmirkhan@yahoo.com)**Abstract**

Oxidative stability of iron fortified butter oil was measured by using different concentrations of anti-oxidants (sesame oil and turmeric powder) in butter oil. The butter oil was prepared by augmented of natural anti-oxidants. i.e. T<sub>0</sub> (100% butter oil+ 30 mg iron sulfate), T<sub>1</sub> (butter oil + sesame oil 5% and turmeric powder 0.10%), T<sub>2</sub> (butter oil + sesame oil 10% and turmeric powder 0.15%), T<sub>3</sub> (butter oil + sesame oil 15% and turmeric powder 0.20%). Samples were stored for 90 days at 40 °C. Butter oil samples were analyzed for free fatty acids, peroxide value, thiobarbituric acid value, shaal oven test and sensorial attributes (color, smell, appearance and overall acceptability) at 30 days of intervals. Treatment T<sub>3</sub> had lowest peroxide and free fatty acid values, while highest values were observed in T<sub>0</sub> and T<sub>1</sub>. Smell, appearance and overall acceptability of iron fortified butter oil shows increased as amount of augmentation of natural anti-oxidants increased in all treatments. The lowest sensory score was observed in treatment T<sub>3</sub> due to dark yellowish type color of turmeric powder. Over all treatments T<sub>2</sub> and T<sub>3</sub> received higher scores from all the physicochemical tests and sensorial evaluation.

**Keywords:** Butter oil, Sesame oil, Turmeric powder, Natural Anti-oxidant.

## STRENGTHENING COMMUNITY RADIO FOR EFFECTIVE CLIMATE CHANGE ADAPTATION STRATEGIES FOR CLIMATE SMART AGRICULTURAL PRODUCTION IN AFRICA

Wilson Okaka<sup>1</sup> and Jennifer Apil<sup>2</sup>

<sup>1</sup>Kyambogo University, <sup>2</sup>Makerere University (Uganda)

[nupap2000@yahoo.com](mailto:nupap2000@yahoo.com)

### Abstract

Strengthening community radio (CR) is crucial for climate change adaptation and mitigation policy awareness of the risk and opportunities for improved potato production in Africa. CR can disseminate climate change adaptation and mitigation information on the new opportunities for agriculture. The principal objectives are to: explain the role of community radio in promoting potato in the food and nutrition policy; state food and nutrition systems policy issues; discuss gender equality and women empowerment for sustainable agriculture and food policies; and explain effectiveness of CR in achieving sustainable development goals two and thirteen (SDGs 2, 13). This policy review employed Ugandan community radio (CR), climate change, food and nutrition policies and best practice for agriculture and food systems information diffusion to achieve SDGs 2, 13. In the searches, we looked for data on enhancing the role of community radio for climate change adaptation and mitigation policy awareness of the risk and opportunities for improved potato production in Africa. The data were then examined for evidence. The retrieved data were summarized for this paper. Community radio is vital for early adoption and diffusion of agricultural technology information for achievement of the SDGs with a focus on SDG 2, and SDG 13. CR radio provides the best coordination for community media stakeholders for agriculture, food, and nutrition research, education, technology, innovations, and policy communication. Community radio services are vital but local radio stations are weak; lack capacity or content, good investments, equipment, well trained staff, and suffer political abuses. Community radio capacity can be well developed to promote climate change smart agricultural opportunities for sustainable livelihoods with a focus on technology innovations diffusion. Africa needs enhanced research funding, policy, capacity building, networking, innovations, South – South and North – South partnerships urgently.

**Key words:** Africa, climate change, climate smart agriculture, radio, technology, diffusion, SDGs

**PUBLIC AWARENESS COMMUNICATION CAMPAIGNS FOR DIFFUSION OF FOOD AND NUTRITION SECURITY POLICY INFORMATION SERVICES FOR SUSTAINABLE LIVELIHOODS IN UGANDA**

Wilson Okaka<sup>1</sup> and Jennifer Apil<sup>2</sup>

<sup>1</sup>Kyambogo University, <sup>2</sup>Makerere University (Uganda)  
nupap2000@yahoo.com, jeinaguma@gmail.com

**Abstract**

Public awareness communication campaigns can end basic hunger, achieve food security and improved nutrition for sustainable agriculture in Uganda. The key objectives are to: describe public communication campaign strategies for sustainable development goal two (SDG 2); discuss the areas of focus in achieving food and nutrition security; highlight the major challenges and prospects of good governance in policy innovations information diffusion and discuss the social responsibility of the press in promoting good governance for food and nutrition policy. In this review, published evidence on the current status of agriculture, food security, and nutrition policy and research innovations diffusion in Uganda was obtained. Information was accessed using internet search engines and libraries. The data that were obtained during the process were used to broaden the search for primary information sources. The findings were contextualized according to the national agricultural, food, and nutrition policy. The retrieved data were summed up for this paper. Public communication campaigns are vital for promoting food and agriculture policy and programme information dissemination for better nutrition in Africa. Effective communication campaigns promote rapid adoption and diffusion of agriculture, health, food, and nutrition technology policy innovations in society. The campaign should be participatory, sustainable, theory driven, gender sensitive, and regularly evaluated for better performance. There are some good north-south, south-south, and inter-university research partnerships which should be consolidated and sustained. Most policy research messages do not reach the target public. The media is the most efficient and effective means of speeding up agriculture, food, and nutrition policy information diffusion. There is urgent need for enhanced north-south collaborative policy research, training, capacity building, technology transfer, innovations funding, ethical standards, good governance, community media development, networking, and gender equality in Uganda.

Key words: Uganda, SDGS, media, food security, agriculture, policy, north-south collaborations



**AN INVESTIGATION ON CHEMICAL COMPOSITION AND NUTRITIONAL EVALUATIONS OF SOME NATURALLY PRODUCE FISHERIES RESOURCES OF PAKISTAN**

\*Zubia Masood<sup>1</sup>, Nelofer Jamil<sup>2</sup>, Sarzamin Khan<sup>3</sup>

<sup>1</sup>Department of Zoology, Jinnah University for Women, Karachi, Pakistan; <sup>2</sup>Department of Chemistry, Sardar Bhadr Khan Women University, Quetta, Pakistan; <sup>3</sup>Department of Poultry Science, Faculty of Animal Husbandry and Veterinary Sciences, The University of Agriculture, Peshawar, Khyber Pakhtunkhwa-Pakistan

[zubiamasood12@gmail.com](mailto:zubiamasood12@gmail.com)

**Abstract**

Hunger and malnutrition are now the most demoralizing issues and are inextricably interconnected with poverty in under developing countries. Today, about one in eight people were found to be suffering from chronic hunger. But nowadays, it has been widely approved around the world that fisheries resources have great capacity to address these challenges and could contribute categorically towards the suppression of hunger, food insecurity and malnutrition. Therefore, the role fish food in providing both adequate food resources and nutrition security for all has now been recognized at universal level. A study was conducted to determine the chemical composition found in the scales, shells and muscle tissues obtained from some commercially important fishes and shellfishes of Pakistan coast in order to observe their nutritional importance for human. Samples were examined for detecting the contents of crude protein, crude lipids, amino acids and certain minerals. Amongst all other animal resources, all these fisheries resources were found to be enriched in protein, certain types of useful amino acids and minerals. Thence, it had been concluded that all these fisheries resources can also be used as useful food resources and bio-products in the productions of useful materials in pharmaceuticals and cosmetic industries.

**Keywords:** Fishes, shellfishes, bio-products, economic values.

## EPIDEMIOLOGY AND CONTROL OF GASTROINTESTINAL FAUNA OF EUROPEAN ALBINO RABBITS OF FAISALABAD DISTRICT, PUNJAB, PAKISTAN

Muhammad Sohail Sajid<sup>\*1,2</sup>, Waqas Altaf<sup>1</sup>, Muhammad Kasib Khan<sup>1</sup>, Muhammad Saqib<sup>3</sup>

<sup>1</sup>Department of Parasitology, University of Agriculture, Faisalabad, Pakistan; <sup>2</sup>U.S. Pakistan Center for Advanced Studies in Agriculture and Food Security (USPCAS-AFS) University of Agriculture, Faisalabad, Pakistan; <sup>3</sup>Department of Clinical Medicine and Surgery, Faculty of Veterinary Science, University of Agriculture, Faisalabad, Pakistan  
[drsohailuaf@hotmail.com](mailto:drsohailuaf@hotmail.com)

### ABSTRACT

Gastrointestinal (GI) Parasitism is one of the main constraints limiting the production of livestock population including rabbits. Different antihelmintics have been used by veterinary practitioners and farmers; however, unable to get rid of GIT parasitism due to multiple factors. This study was aimed to determine the frequency distribution and control of GI parasites of rabbit population of Faisalabad district. To this end, a questionnaire was developed to record the epidemiological parameters which associated with the disease distribution in rabbits. Fresh faecal sample were collected for lab analysis for screening the helminth positive subjects. Positive rabbits for worms were subjected to oral formulations of levamisole (LEV), albendazol (AB), ivermectin (IVM) and Pyrental pamoate (PP). Total 384 rabbits were screened from five study sites of the selected district. Overall prevalence of Gastrointestinal Parasitism in rabbit Populations of study district was found 84.37 %. Eggs of Trichostrongylus retortaeformis, Graphidium strongsum and Cittotenia ptenoids were seen in feecal egg count assay 88.74%, 8.74% and 2.54% respectively. In in vivo drug trials the efficacy of IVM was good as compared to all other drugs. Efficacy of LEV was satisfactory but as compared to other drugs not remarkable. Effect of PP and AB were almost same. So, IVM is recommended for GI parasitism. The study provided the better choice of drug for preventive and/or therapeutic management of GI parasitism in rabbit population of Faisalabad, Punjab, Pakistan.

Keywords: GI Parasitism, Albino Rabbit, Epidemiology, in vivo, ivermectin

**COMPARATIVE EVALUATION OF MICROBIOLOGICAL MILK SAFETY STANDARDS EXISTING IN FAISALABAD**

Nabila Yaqoob,<sup>1</sup> Ahsan Naveed\*<sup>1</sup>, Sajjad Ur Rahman<sup>1</sup>, Syeda Wajiha Hassan<sup>1</sup>, M. Farooque Hassan<sup>2</sup>

<sup>1</sup>University of Agriculture Faisalabad<sup>1</sup>; <sup>2</sup>Shaheed Benazir Bhutto University of Veterinary and animal sciences Sakrand Sindh, Pakistan

[ahsannaveed.uaf@gmail.com](mailto:ahsannaveed.uaf@gmail.com)

**Abstract**

Milk is highly perishable and easily spoiled by microbial contamination. Pathogenic bacterial contaminants in milk may causes different diseases like dysentery, tuberculosis, salmonellosis and fever. Microbiological quality milk safety standards were evaluated. Milk samples (Raw, Pasteurized, UHT) were collected from farms and commercial market around Faisalabad district and analyzed by different methods like, methylene blue reduction test (MBRT), swab test on animal food (STAF) to check for the antibiotic residues in raw milk, somatic cell count (SCC), total coliform count (TCC), thermoduric count, detection of *M. tuberculosis* and pathogenic *E. coli*. In raw milk samples only 2% were positive for STAF while 5% for pathogenic *E. coli*. The mean SCC, TCC and thermoduric count for raw milk was  $1.5 \times 10^5$ , 298 and 900 respectively. TCC for pasteurized was 150 and it was zero for UHT. The thermoduric count for pasteurized and UHT was 1100 and 1480 respectively. All the three categories were negative for detection of *M. tuberculosis*. Overall results indicated that about 10% of raw and pasteurized milk samples were positive for different type of contaminations and do not meet the milk safety standards.

**Key words:** UHT, pasteurized, STAF, TCC, contaminations, thermoduric

**HIGH OMEGA-6/OMEGA-3 RATIO IN MARGARINE INDUCES INTESTINAL INFLAMMATION AND OXIDATIVE STRESS RESPONSE IN WISTER RAT**

Aisha Mahmood \*, Muhammad Naeem Faisal, Tanweer Khaliq, Junaid Ali Khan, Faqir Muhammad, Humaira Muzaffar, Farooq Azam and Maha Saleem

Institute of Pharmacy, Physiology, and Pharmacology, University of Agriculture  
Faisalabad, Pakistan

[\\*2007ag1551@uaf.edu.pk](mailto:*2007ag1551@uaf.edu.pk)

**Abstract**

Diets have a great impact on health and health-related issues. High-fat diets have been linked to epidemic development of metabolic disorder. Manifestation of metabolic disorder is causally associated with chronic diseases such as insulin resistance, inflammatory bowel disease (Ulcerative colitis and Crohn's disease), intestinal junction disruption and ultimately cancer. Underlying mechanisms are association with oxidative stress. This study was designed to investigate the development of metabolic disorder after a high-fat diet especially with high omega-6/omega-3 ratio in margarine on various physiological aspects, gut pathophysiology, epithelial signature and oxidative stress in a rat model. Results indicated a complex interplay of high ratio of omega-6/omega-3 in high-fat diet-induced oxidative stress on gut epithelium alterations, lipid metabolism, and plasma insulin level. Overall results suggest that high-fat diet as pervasive factors that exacerbate its lethal effect on gut physiology and metabolism. The data were statistically analyzed by one-way ANOVA and DMR.

**Keywords:** Metabolic disorders, gut pathophysiology, metabolic stress, lipid metabolism.

## EFFECT OF GAMMA-IRRADIATION ON THE QUALITY CHARACTERISTICS OF BROWN RICE BASED WEANING FOOD

Awsi Jan, Dr. Monika Sood, P.R.Hussain, Lubna Masoodi, Ambreena, Nadia Bashir  
Shere-e-Kashmir University of Agricultural Sciences and Technology, Jammu and  
Kashmir  
[awsijan@gmail.com](mailto:awsijan@gmail.com)

### Abstract

Scientifically, it has been proved that breast milk is the perfect food for the infant during the first six months after birth. It contains all the nutrients and immunological factors an infant requires to maintain optimal health and growth. Furthermore, breast milk also protects infants against the two leading causes of infant mortality- upper respiratory infections and diarrhoea. On the other hand, nowadays, due to the reduced consumption of breast milk, important nutrients such as proteins, zinc, iron and B-vitamins are likely to be deficient in the contemporary diet of the affected infants, at the age of six months and above when the child's birth weight is expected to have doubled. Hence, breast milk is no longer sufficient to meet the nutritional needs of the growing infant. Since there are many weaning foods available in the market that are processed by conventional thermal methods. This leads to loss in the nutritional as well as the sensory profile of the product. So this work is mainly concerned with the 'Effect of g-irradiation quality characteristics of brown rice based weaning food'. In food technology, recent research has elucidated new potential applications for radiation. High doses of ionizing radiation have been shown to inhibit growth of microbes and to reduce the level of anti nutrients which were reported to reduce the availability of protein and minerals. Gamma Irradiation also results in modification of physico chemical properties like increase in the solubility and decrease in swelling index which has positive influence in the quality of weaning food.

**Key words:** weaning food, brown rice, gamma- irradiation

**PREVALENCE OF ESCHERICHIA COLI O157: H7 IN BEEF MEAT IN WESTERN ALGERIA**

Barka Mohammed Salih and Benyoub Noreddine

Route de la Rocade BP 119 faculté CNV-STU Université de Tlemcen Tlemcen 13000  
Algérie  
[bm\\_salih@hotmail.com](mailto:bm_salih@hotmail.com)

**Abstract**

Actually, there are no official regulations stipulating the procedures for sampling and research of STEC in food in Algeria. In this study, we were interested in search of *E. coli* O157: H7 Shiga-toxin producing (STEC) in 750 samples of beef meat imported from different countries. After enrichment and use of selective agents that are intended to curb the growth of the annex flora, we have isolated 5 strains from meat. These strains presented the main characteristics of *E. coli* O157: H7, non sorbitol fermenting and negative  $\beta$ -Dglucuronidase. Genetic characterization revealed the presence of genes and toxicity *stx1 stx2* responsible for pathogenicity of these bacteria by the production of toxins as well as the specific hemolysin (*ehxA*) and intimin (*eae* genes) for *E. coli* O157: H7. The study of the resistance of the strains to antibiotics disclosed they are sensitive to antibiotics tested.

**Keywords:** *E. coli* O157: H7, beef meat, *stx1*, *stx2*, antibiotics

**RISK CHARACTERIZATION OF ANTIBIOTIC RESISTANCE OF SALMONELLA ISOLATED FROM RAW MEAT IN PENANG, MALAYSIA**

Hafiz NidaUllah, Mohd Omar Abdul Kadir, Rosma Ahmad, Widad Fadhullah and Saima Sohni

School of Industrial Technology, Universiti Sains Malaysia, 11800, Georgetown, Penang MALAYSIA

[hafiz.nidaullah@daad-alumni.de](mailto:hafiz.nidaullah@daad-alumni.de)

**Abstract**

This research work has been endeavoured to estimate the qualitative risk characterization of antibiotic resistance and genetic determinants of resistance of Salmonella isolated from raw meat samples. A total of 400 samples including beef, chicken, mutton and fish meat from different wet markets (240) and hypermarkets (160) in Penang, Malaysia, were collected. Detection of Salmonella was carried out by following ISO (6579: 2002), susceptibility against 15 antibiotics of clinical importance and 13 different resistance genes in Salmonella were investigated in this study. Further, risks of antibiotic resistance to consumers were investigated using risk analysis approaches. As a result, Salmonella expressed maximum resistance to two antibiotics; tetracycline and sulfamethoxazole from beef, chicken, mutton and fish in wet markets and to sulfamethoxazole from beef, chicken and fish meat in hypermarkets. Highest prevalence of resistant genes in Salmonella was found for streptomycin (strpA, strpB= 80%) and tetracyclin (tetA=80%) in wet markets and streptomycin (strpA, strpB= 80%), amoxicillin (bla-TEM: 80%) and sulfamethoxazole (Sul12: 100%) in hypermarkets. Moreover, risk characterization model signified four cases of Very High Additional Risk (VHAR) of resistance; three cases of streptomycin from beef, chicken, fish and one case of ampicillin from fish. In conclusion, increasing antibiotic resistance, due to the potential prevalence of antibiotic resistant genes in Salmonella and ultimately the originated risk of resistance against WHO enlisted life saving antibiotics highlights the emergence of antimicrobial resistance in food chain.

**Keywords:** Raw meat, Salmonella, Antibiotic resistance

## COMPARATIVE EVALUATION OF NIGELLA SATIVA AND LINCOMYCIN AS GROWTH PROMOTERS IN COMMERCIAL BROILER

Munnaza Shareef, Muhammad Naeem Faisal, Aisha Mahmood, Humaira Muzaffar,  
\*JazibHussain, Samia Ali, Muhammad Jamal, Farooq Azam.

Institute of Pharmacy, Physiology & Pharmacology, University of Agriculture,  
Faisalabad, Pakistan

[jazibhussain04@gmail.com](mailto:jazibhussain04@gmail.com)

### Abstract

*Nigella sativa* (*N. sativa*) is an important herb with multiple pharmacological activities. Therapeutic properties of *N. sativa* are based on the availability of thymoquinone (TQ). Current study investigates the antibacterial effect of *N. sativa* in comparison to lincomycin as growth promoter in broiler. Effect of *N. sativa* powder on feed conversion ratio (FCR) was also observed in broiler chicks. On weekly basis body weight of birds was observed. Immune titre against Newcastle and Infectious Bursal disease was also determined. Statistically analysis of variance and significance ( $P < 0.05$ ) of the results was determined through Duncan's Multiple Range test. Result demonstrated that supplementing the broiler feed with 1% *N. sativa* improved body weight gain and feed conversion ratio of broilers at finisher phase ( $P < 0.05$ ). In conclusion, results have shown that addition of 1% *N. sativa* powder appear to have a positive impact on growth performance of broiler chicks and it could be considered as a substitute of antibiotic growth promoter (lincomycin) for broiler chicks.

**Key words:** *Nigella sativa*, lincomycin, Immune titer, FCR, growth promoter, histopathological



**ESTIMATION OF MILK PROTEASE ACTIVITY FROM UNINFECTED MAMMARY GLANDS OF NILI-RAVI BUFFALOS , SAHIWAL AND CROSS-BREED COWS OF PAKISTAN.**

Razia Kausar<sup>1</sup>, Amjad Hameed<sup>1</sup>, and Zafar Iqbal Qureshi<sup>2</sup>

Department of Anatomy, FVS, University of Agriculture, Faisalabad, Paksitan; Nuclear Institute of Agriculture and Biology<sup>1</sup>, Jhang Road Faisalabad, Pakistan; Department of Theriogenology; <sup>2</sup>Faculty of Veterinary Science, The University of Agriculture, Faisalabad, Pakistan

[razia\\_uaf@hotmail.com](mailto:razia_uaf@hotmail.com)

**Abstract**

Pakistan is at 4th place among largest milk producing countries of the world and it produces 49.5 million tons of milk annually. Dairy industry of Pakistan consists of buffaloes and cows. The share in annual milk production of country by cow is 17.37 million tons and by buffalo is 30.46 million tons. The level of protease activity in milk was tested and compared in milk samples from Nili-Ravi buffaloes, Sahiwal and cross-bred cows. The milk protease activity differed significantly ( $p < 0.05$ ) among tested dairy species/cattle. The mean value for protease activity in milk was highest for cross-bred cows followed by Sahiwal cows and lowest for Nili-Ravi buffaloes. With a minimum value of 32 units/mL and a maximum value of 95 units/mL, the mean value of milk protease activity for Nili-Ravi buffaloes was  $57.3 \pm 2.16$  units/mL. For Sahiwal cows, the minimum value of milk protease activity was 33 units/mL while maximum value was 99 units/mL with a mean value of  $69.6 \pm 3.02$  units/mL. For cross-bred cows, minimum value of milk protease activity was 39 units/mL and maximum value was 127 units/mL with a mean value of  $81.3 \pm 3.35$  units/mL. As for as protease activity was concerned, the milk of Nili-Ravi buffaloes was comparatively better as compared to other two dairy animal types investigated. Moreover, values being reported can be used as reference values of protease activity in milk for Nili-Ravi buffaloes, Sahiwal and cross-bred cows.

Keywords: Pakistan, milk, Nili-Ravi

**APPLICATIONS OF PULSE ELECTRIC FIELD IN DAIRY TECHNOLOGY**

Saleem Malik, Lubna Masoodi , Awsi Jan, Saleem Malik

Sher-e-Kashmir University of Agricultural Sciences and Technology Kashmir Srinagar-  
Shalimar- 190025[malik.saleem72@gmail.com](mailto:malik.saleem72@gmail.com)**Abstract**

Thermal processes used for the preservation alter the nutritional and sensory attributes of food materials. The food industry is aiming for new preservation techniques to replace the thermal processes with non- thermal processes, where there is no generation of heat thus have minimal effects on nutritional and sensory attributes of food. The thermal process pasteurization which is used to extend the shelf life of milk and inactivates the pathogenic bacteria, affect the nutritional and organoleptic properties of milk. The main effect on milk is the development of cooked flavor. In order to overcome such causes pulse electric field is used. Pulse electric field is considered as one of the most promising non-thermal processing technology. The pulse electric field (PEF) is alternative method to thermal pasteurization for milk that have minimal effects on sensory and nutritional attributes of milk thus providing fresh like products. Most PEF systems used for treatment of dairy or dairy products have been limited to bench top or small pilot scale systems. The PEF technology used in milk inactivates the microorganisms and enzymes and it has been reported that PEF have minimum adverse effects on sensory attributes of milk. It has been found that PEF treatment (35 kV/cm; 2.3  $\mu$ s of pulse width at 65<sup>0</sup>C for < 10s) immediately given to milk after pasteurization extended the shelf life of milk to 78 days at 4<sup>0</sup>C. PEF technology is considered as a promising technology in dairy industry. Its major application is on inactivation of vegetative microorganisms. It has least effects on color, taste and nutritive value of dairy and dairy products. It is effective in extending the shelf life of milk.

**Key words:** Dairy, Dairy products, PEF, Pasteurization.

**EFFECT OF MENTHA PIPERITA (PEPPERMINT) EXTRACT AND ITS JUICE ON EGG QUALITY TRAITS DURING DIFFERENT STORAGE TIME IN LAYING HENS**

Abdur Rahman<sup>1</sup>, E. Eren Gültepe<sup>2</sup>, Cangir Uyarlar<sup>2</sup>, I. Sadi Çetingül<sup>2</sup>, Aamir Iqbal<sup>1</sup>, Ismail Bayram<sup>2</sup>

Aamir Iqbal

<sup>1</sup>Department of Animal Nutrition, University of Veterinary and Animal Sciences, Lahore-Pakistan

<sup>2</sup>Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, Afyon Kocatepe University, Afyonkarahisar, Turkey.

aamir\_vet@yahoo.com

**Abstract**

Much focus has been given on the use of herbs and herbal products to improve performance and to some extent the quality in freshly laid eggs but limited data are available regarding the impact of herbs on storage quality of eggs. The present study was designed to evaluate the effect of *Mentha Piperita* oil and mentha juice in feed and drinking water respectively, on egg quality traits in laying hens at different storage intervals. A total of 252 Babcock laying hens were divided into 7 groups and each group was further divided into 4 subgroups having 9 hens in each. Group A served as a control. Group A was fed basal diet without any supplementation. Group B, C and D were offered diets supplemented with mentha extract 50, 100 and 200 mg/kg of feed while groups E, F and G diets were having same doses of mentha juice in drinking water. At the end of the study (56 days), a total of 252 eggs (36 eggs from each group) were collected randomly. 84 eggs were analyzed at zero day of storage while other eggs were stored at 4°C temperature. Among these eggs, 84 were analyzed after 15 days and remaining 84 after 30 days of storage. The results revealed that egg quality traits like egg shell breaking strength (ESBS), yolk color (YC), haugh unit (HU) and egg weight showed non-significant difference ( $P>0.05$ ) among all the groups at different storage intervals.

**Key Words:** Egg Quality, Mentha piperita, Haugh unit, Yolk color, Egg shell breaking strength

**THE RECENT DEVELOPMENTS IN PREVENTING EMBRYONIC LOSSES IN SMALL RUMINANTS DURING HEAT STRESS.**

Dehouegnon Jerry Agossou, Uğur Serbest, Nazan Koluman  
Department of Animal Science, Çukurova University, Turkey  
bermondjerry@gmail.com

**Abstract**

Exposure to elevated ambient temperature during period prior to conception and the early stages of pregnancy has strong adverse effects on reproductive outcomes such as reducing the quantity and quality of ovine and goat embryos. An inappropriate feeding (overfeeding or underfeeding) of reproductive does and ewes affects blood progesterone concentrations and metabolic hormones such as Insulin-like Growth Factor I (IGF-I) altering oocyte and embryo quality. On the other hand, the deficiency of some vitamins and minerals especially vitamin E and selenium can cause accumulation of lipid peroxidation products which can lead to preeclampsia or pregnancy-induced hypertension. However, the housing practices and well feeding by supplying balanced dietary energy and polyunsaturated fatty acids (PUFAs) pre-and during mating may improve embryo survival. Studies suggest that hormonal treatment with gonadotropin-releasing hormone (GnRH) fluorogestone acetate (FGA), arginine may reduce embryo mortality. This article aims to describe factors affecting embryo losses and explore some of the recent developments in prevention early embryo losses in small ruminants.

**Keywords:** heat stress, pregnancy, sheep, goat

**COMPARISON OF LINCOMYCIN AND TETRACYCLINE AS ANTIBIOTIC GROWTH PROMOTER IN COMMERCIAL BROILER**

Muhammad Jamal, Muhammad Naeem Faisal, Hamza Nazeer, Humaira Muzaffar, Aisha Mahmood, Jazib Hussain, Jawad Aslam, Hira Javed

Hamza Nazeer

Institute of Pharmacy, Physiology and Pharmacology, University of Agriculture, Faisalabad-Pakistan

[hamzanazir56@yahoo.com](mailto:hamzanazir56@yahoo.com)

**Abstract**

Antibacterial agents are the most valuable means available against bacterial infections such as, lincomycin and tetracycline are common brand. It is a cost-effective drug because low cost, good activity against anaerobic bacteria, favourable pharmacokinetic and pharmacodynamics. Still the criterion standard for therapy of anaerobic infections was especially in gut infection. Use of sub therapeutic level of antibiotic in animal feed as growth promoters are primarily source of antibiotic resistance and stress factor for gut epithelium. Current study was designed to elaborate the advances in understanding comparison in between lincomycin and tetracycline as AGPs in commercial broiler. Physical parameters, immune titer (Newcastle and Infectious Bursal disease), histopathological examination of gut and hematological parameters was studied. At the same time, feed intake (g) and feed conversion ratio was measured accordingly for hematological and histopathological analysis. Results of this study show that addition of tetracycline increases immune titer against ND and IBD diseases as compared to control and lincomycin treated groups. Results showed that lincomycin has significantly perform better as compared to tetracycline while as, tetracycline showed resistance against bacterial population. Statistically data was analyzing through analysis of variance and significance of the result was determined through Duncan's Multiple Range test.

**Key Words:** IBD, ND, Feed conversion ratio (FCR), AGPs, Tetracycline

## PHYTOREMEDIATION: AN INSPIRING AND ECONOMICAL FOR THE TREATMENT OF IRRIGATION WATER

Hina Jabeen, Aqsa Gilani, Fatima Waleed, Mushtaq Ahmad, Rakhshinda Sadiq, Akhtar Rasool

Women University Mardan, KPK Pakistan

[hinajabeen2@gmail.com](mailto:hinajabeen2@gmail.com)

### Abstract

Wastewater irrigation is a common practice in developing countries. This practice poses a high risk to food security and human health due to contaminants of emerging concern (CECs) contained in the wastewater. These CECs originate from industrial effluents and domestic discharges which contain persistent organic pollutants (POPs), residues of pesticides, insecticides, dyes and pharmaceutical drugs etc. These toxicants are taken up by the crops making way to enter the food chain, bio-accumulate at all trophic levels and present a terrific risk to food security. In this scenario, remediation of wastewater is becoming a big challenge worldwide. Conventional methods are very costly and environment unfriendly due to noxious by-products. Therefore, phytoremediation, an innovative field of environmental bio-technology, is becoming an ultimate alternate of the conventional methods. Phytoremediation exploits the natural detoxification potential of plants alongwith their associated microbes to remediate polluted water. This study presents a remarkable potential of two aquatic plants, *Nasturtium officinale* (watercress) and *Lemna minor* (duckweed) to remediate domestic wastewater of Shehzad town, Islamabad, Pakistan. The study of physico-chemical parameters of wastewater showed very high levels of chemical and biological oxygen demand (COD, BOD), total dissolved solids (TDS), total hardness, pH and turbidity. Interestingly, following the treatment with the designated plant system, a significant decline was observed in all the studied parameters which were comparable to irrigation water standards. Hence, the two plants presented the marvellous capability to extract and degrade the pollutants from wastewater which was obvious from the colour and odour of water. Henceforth, the proposed technology would serve as a great asset in the already existing repertoire of efforts to produce pollutants free food. Moreover, good irrigation practices will provide an excellent step towards food security by eradicating the poisons from food crops.

**Keywords:** Wastewater, irrigation, human health, contamination

**COMPARATIVE EVALUATION OF NIGELLA SATIVA AND LINCOMYCIN AS GROWTH PROMOTERS IN COMMERCIAL BROILER**

Munnaza Shareef, Muhammad Naeem Faisal, Aisha Mahmood, Humaira Muzaffar, \*JazibHussain, Samia Ali, Muhammad Jamal, Farooq Azam.

Institute of Pharmacy, Physiology & Pharmacology, University of Agriculture, Faisalabad, Pakistan

[jazibhussain04@gmail.com](mailto:jazibhussain04@gmail.com)

**Abstract**

*Nigella sativa* (*N. sativa*) is an important herb with multiple pharmacological activities. Therapeutic properties of *N. sativa* are based on the availability of thymoquinone (TQ). Current study investigates the antibacterial effect of *N. sativa* in comparison to lincomycin as growth promoter in broiler. Effect of *N. sativa* powder on feed conversion ratio (FCR) was also observed in broiler chicks. On weekly basis body weight of birds was observed. Immune titre against Newcastle and Infectious Bursal disease was also determined. Statistically analysis of variance and significance ( $P < 0.05$ ) of the results was determined through Duncan's Multiple Range test. Result demonstrated that supplementing the broiler feed with 1% *N. sativa* improved body weight gain and feed conversion ratio of broilers at finisher phase ( $P < 0.05$ ). In conclusion, results have shown that addition of 1% *N. sativa* powder appear to have a positive impact on growth performance of broiler chicks and it could be considered as a substitute of antibiotic growth promoter (lincomycin) for broiler chicks.

**Key words:** *Nigella sativa*, lincomycin, Immune titer, FCR, growth promoter, histopathological

**EFFECT OF ANTIOXIDANT, PACKAGING MATERIAL AND STORAGE PERIODS ON THE CHEMICAL COMPOSITION OF ALMOND KERNEL**

Lubna masoodi, Dr A.H Rather, Awsi Jan  
Shere-e-Kashmir University of Agricultural Sciences and Technology Srinagar. 190025  
Jammu and Kashmir  
[lubna.masudi@gmail.com](mailto:lubna.masudi@gmail.com)

**Abstract**

The almonds of uniform size were manually decorticated and extracted kernels were mechanically dried at 40<sup>0</sup>C to final moisture content of 4.20% after antioxidant treatment (BHA+BHT @ 0.015%) were given to each. The treated samples were packed in LDPE and laminates (vaccum) to monitor the changes in quality attributes under ambient and accelerated (35<sup>0</sup>C) storage conditions for a period of 270 days. Antioxidant treated samples packed in vaccum laminates and stored under ambient conditions proved significantly superior by exhibiting highest crude fat content (54.33%) crude protein content (15.16%) crude fiber content (9.81) minimum peroxide value (109.99 meq/100g) and free fatty acid value (0.95mg/g). compared to the untreated kernels packed in LDPE under accelerated conditions throughout the storage periods. Results of present study have led to the conclusion that antioxidant treatment vaccum packaging and ambient conditions proved best in maintaining the quality of almond during storage.

Key words: Almonds antioxidants and packaging material



**AN ANALYSIS OF CORRELATION BETWEEN POOR FOOD EDUCATION AND DENTAL CARIES IN PESHAWAR**

Maria Nizam

Sardar Begum Dental College, Gandhara University Peshawar

[Marianizam65@gmail.com](mailto:Marianizam65@gmail.com)

**Abstract**

Poor food education leads to intake of refined carbohydrates such as candies, cookies, cakes, soft drinks, soda, honey, processed foods such as the fries, pasta, biscuits, pre-sweetened cereals and low quality food that in turn causes a high risk factor in development of caries. Data on quantification of poor food education and carries is deficient in medical and dental literature in context to Pakistan. Hence an analytical cross sectional study to determine the prevalence of caries among adolescents of age 16-18 years was done in Peshawar. Total sample size was 100 adolescents. Data was collected through a self-administered questionnaire. The study was carried out to determine the causes of prevalence dental caries among adolescents in Peshawar, to assess the knowledge of oral health among adolescents in Peshawar and to quantify the data on food education and dental caries among adolescents in Peshawar. The collected data suggests that poor Food Education is positively correlated with Dental Caries.

**Keywords:** Food Education, Dental Caries, Carbohydrate Rich Food, Peshawar

**MONOGLYCERIDES AN ALTERNATIVE TO ANTIBIOTIC GROWTH PROMOTERS IN COMMERCIAL BROILER**

Samia Ali, Muhammad Naeem Faisal, Aisha Mahmood, Humaira Muzaffar, Muhammad Jawad Aslam\*

Institute of Pharmacy, Physiology & Pharmacology, University of Agriculture, Faisalabad-Pakistan

[jawadaslamch@gmail.com](mailto:jawadaslamch@gmail.com)

**Abstract**

A wide use of antibiotic growth promoters (AGPs) is beneficial and helpful for increasing efficiency which in turn makes animals healthy and more productive. The AGPs have also their role in reducing harmful bacteria from gut and reducing the competition for host nutrients as well as reduced the subclinical disease caused by these bacteria to obtain better performance. Current study investigates the antibacterial effect of Monoglyceride (Monolaurin) in comparison to lincomycin as growth promoter in broiler chicks. Effect of Monoglycerides on feed conversion ratio, physical parameters, immune titer (Newcastle and Infectious Bursal disease), histopathological examination of gut, screening antibacterial activity were performed. Result showed that supplementing the broiler feed with Monoglyceride (monolaurin) improved body weight gain and feed conversion ratio (FCR) of broilers at finisher phase. Results showed that addition of Monoglyceride (Fatty acid) appear to have a positive impact on growth performance of broiler chicks and it could be considered as a substitute of antibiotic growth promoter (lincomycin). Statistical data was analyzed by applying ANOVA and DMR. In conclusion, Monoglycerides showed better results as compared to lincomycin on the basis of better FCR, Immune titer, and gut health in commercial broiler.

**Key words:** Monoglycerides, lincomycin, Immune titer, FCR, growth promoter, histopathological examination

## EFFECT OF FENUGREEK SEED SUPPLEMENTATION ON EGG QUALITY AND PRODUCTION PERFORMANCE OF SPENT LAYERS

N. Chand., F.Wahab, M.S. Qureshi and R.U. Khan

Faculty of Animal Husbandry & Veterinary Sciences, The University of Agriculture, Peshawar, Pakistan

[draleeze@yahoo.com](mailto:draleeze@yahoo.com)

### Abstract

The present research study was conducted to investigate the effect of various levels of fenugreek seeds supplementation on feed intake, egg production, egg quality and liver health of Rhode Island Red spent layers at the research unit of the University of Agriculture Peshawar, Pakistan. A total of 60 spent Rhode Island Red layers were assigned to four groups (FS-0, FS-0.5, FS-0.75, and FS-1) with three replicate having 5 birds each and were kept in cages for 28 days in open sided house. Group FS-0 was kept as control while groups FS-0.5, FS-0.75, FS-1 were provided with fenugreek seed powder in the ration at the rate of 0.5, 0.75 and 1% respectively. Egg production and feed intake was recorded on daily basis and from daily data weekly and over all egg production and feed intake was determined. Egg shell weight, egg weight, egg shell thickness, Haugh unit, albumin height and yolk weight were determined on weekly basis. Dietary supplementation of fenugreek seed significantly ( $P < 0.05$ ) decreased feed intake at all recorded stages. At first week significantly higher feed intake was recorded for the control group ( $105.06 \pm 0.279$ ) and was followed by group FS-0.5 ( $103.79 \pm 0.067$ ), FS-0.75 ( $103.57 \pm 0.058$ ) while lowest intake of feed was noted for group FS-1 ( $101.59 \pm 0.278$ ). The same trend was observed during week-2, -3 and -4. Overall higher feed intake was noted for the control group ( $105.16 \pm 0.198$ ) while lower intake of feed was noted for group FS-1 ( $101.93 \pm 0.306$ ). Significantly higher ( $P < 0.05$ ) egg production was recorded for group FS- 0.5 followed by group FS-0.75 while lower egg production was recorded for the control and group FS-1. Egg quality traits including egg shell weight, egg weight, egg shell thickness, Haugh unit, albumin height and yolk weight were ( $P > 0.05$ ) not affected by supplementation of fenugreek seeds. Supplementation of fenugreek seeds improved liver function test indicated by decreased liver enzymes including Alanine aminotransferase (ALT) and Aspartate aminotransferase (AST). Significantly higher ( $P < 0.05$ ) ALT and AST levels were recorded for the control while lower ( $P < 0.05$ ) ALT and AST values were recorded in supplemented groups. It is concluded from the present study that supplementation of fenugreek at the rate of 0.5% increased egg production without affecting egg quality traits and liver health.

**Keywords:** Layers, egg quality, cholesterol,

**DETERMINATION OF RED AND WHITE MEAT CONSUMPTIONS OF ADULT INDIVIDUALS LIVING IN AKŞEHİR**

Nazik Meziyet DİLEK, Ezgi TOPTAŞ BIYIKLI, Ali Emrah BIYIKLI  
University, Kadir Yallagöz School of Health, Department of Nutrition and Dietetics  
[meziyetteemel@hotmail.com](mailto:meziyetteemel@hotmail.com)

**Abstract**

This study has been carried out to the red and white meat consumptions of the people in the district Akşehir. This study was carried out on 300 adults (age 18 and over), who reside in Akşehir and are randomly selected, in 2017 January and February. In the study, a questionnaire, formed as a result of literature review, was used as a data collecting instrument. According to the data obtained, 77.3% of those participating in the study are females and 22.7% are males. It was identified that 54.7% of the participants were in the range of ages 18-30, 30.8%, ages 31-50, and 14.5%, age 50 and over. It was found that 49.3% of the participants consumed red meat the most in their diets; 44.00%, poultry meat; and 6.7%, fish and other products. When the consumption amount of monthly red meat of the individuals participating in the study were assessed, it was identified that 42.8% of them consumed red meat less than 500 g; 23.4%, 500 to 1000 g; 19.4%, 1001 to 1500 g; and 14.4%, more than 1500 g. When those participating in the study were assessed in terms of monthly fish meat, it was found that 64.7% of them consumed fish meat less than 500 g and 5%, more than 1500 g. At the end of the study, although the most preferred sort of meat was red meat, it was identified that the large majority (42.8%) of those participating in the survey consumed red meat less than 500 g. Fish was identified as the sort of the least preferred meat in diet, it was concluded that 64.7% of the participants consumed fish meat less than 500 g. The reasons for consuming less meat in our society should be researched and precautions should be developed toward solution.

**Keywords:** Red meat, adult, fish, poultry

## IDENTIFICATION OF THE PREFERENCES OF HEALTHY ADULTS ABOUT RED AND WHITE MEAT CONSUMPTION

Nazik Meziyet DİLEK, Ali Emrah BIYIKLI, Ezgi TOPTAŞ BIYIKLI  
Selcuk University, Kadir Yallagöz School of Health, Department of Nutrition and Dietetics  
[meziyetteemel@hotmail.com](mailto:meziyetteemel@hotmail.com)

### Abstract

This study was carried out to identify the preferences of the healthy adults about red and white meat consumption. This study was carried out on 300 adults, who reside in Akşehir and are randomly selected, in 2017 January and February. In the study, a questionnaire, formed as a result of literature review, was used as a data collecting instrument. It was found that 73.4% of those participating preferred bovine meat; 18.3%, mutton, and 18.3%, goat's meat. The preferences of the participants to purchase red meat were found as meat cubes with 52.9%; mincemeat with 34.6%; fillet steak with 4.2%; chopped steak with 3.1%; and beefsteak with 2.8%. While chicken meat is almost completely (99%) as poultry meat, the rate of those preferring turkey and bird meat was identified to be 0.7% and 0.3%, respectively. The preferences to purchase poultry meats were identified as whole body with 52.7%; chicken breast with 25%; and drumstick with 18%. In the preferences to purchase fish meat, anchovy was identified as the most preferred sort of fish with 58.7% and this was followed by salmon with 25.7%. It was identified that the criteria the individuals participating in the study pay attention the most, while purchasing the red and white meat, as date of expiry (70.7%), color and appearance (19.7%), brand and firm (4%), packing (1.7%), price (1.7%), nutritional value (1.7%), and other factors (0.7%). It was found that the most preferred meat was bovine meat as red meat; chicken meat as poultry meat; and anchovy as fish meat. The criterion the participants pay attention the most, while purchasing the red and white meat, was identified to be date of expiry. With the similar studies to be carried out in the different regions, the preferences of our society can be evaluated more comprehensively.

**Keywords:** Meat, healthy, adults, chicken

**CONJUGATED LINOLEIC ACID IN ANIMAL SOURCE FOODS AND ITS EFFECT ON HEALTH**

Nazik Meziyet DİLEK, Mustafa KARAKAYA

Selcuk University, Kadir Yallagöz School of Health, Department of Nutrition and Dietetics

[meziyetteemel@hotmail.com](mailto:meziyetteemel@hotmail.com)

**Abstract**

Conjugated linoleic acid (CLA) is a positional and geometrical isomer of linoleic acid and the most common isomers 9c,11t-18:2 and 10t,12c-18:2. CLA isomers present the most commonly in the meat, milk, and other products, which are obtained from ruminant animals, while in egg they are relatively less. Synthesizing isomers from linoleic is carried out in two different ways and these are microbial bio-hydrogenation of linoleic acid to stearic acid in rumen or desaturation of trans vaccenic acid to CLA isomers through  $\Delta^9$ -desaturase enzyme in mammalian glands and adipose tissues. That CLA isomers, which has anti-carcinogenic, anti-atherogenic, and anti-diabetic are provided from animal resources such as meat, milk and meat led the necessary studies to be carried out to increase CLA concentrations in these products. In this study, it has been attempted to give information about the sources of CLA and factors influencing CLA content in these sources and effects of CLA on health.

**Keywords:** Fatty acids, meat, milk, poultry

**IDENTIFICATION OF PROCESSED MEAT PRODUCTS CONSUMPTION OF HEALTHY ADULT INDIVIDUALS LIVING IN AKŞEHİR**

Nazik Meziyet DİLEK, Ezgi TOPTAŞ BIYIKLI, Ali Emrah BIYIKLI  
Selcuk University, Kadir Yallagöz School of Health, Department of Nutrition and Dietetics  
[meziyetteemel@hotmail.com](mailto:meziyetteemel@hotmail.com)

**Abstract**

This study was carried out to identify the processed meat products consumption of the people living in the district Akşehir. This study was carried out on 300 adults (age 18 and over), who reside in Akşehir and are randomly selected, in 2017 January and February. In the study, a questionnaire, formed as a result of literature review, was used as a data collecting instrument. According to the data obtained, 61.8% of those participating in the study are females and 38.2% are males. While 17.8% of the participants stated that they found processed meat products healthy, 62.8% expressed that they found it unhealthy and 39.4% did not have any ideas. While 35.5% of the individuals participating in the survey had information about the production of processed meat products, 33.6% expressed that they did not have any information about it and the remaining 30.9% did not have any idea. 36.8% of those participating in the study consumed the processed meat product at least once a week; 26.6%, every fifteen days; and 21.7%, once a month. While 6.9% of those participating consume processed meat product every day, 7.9 % don't consumes processed meat products at all. As a result of the study, although 62.8% of those participating in the survey consider that processed meat products are unhealthy, only 35.5% of them expressed that they had information about the production of these products. Although the large majority of those participating in the study (62.8%) consider that the processed products are unhealthy, only 7.9% do not consume these products. As a result of the study, it was seen that the society did not have enough information about this issue. The reasons for this state should be researched and the society should be adequately informed about this issue.

**Keywords:** Meat, processed, health, information

**POTENTIAL OF RED MEAT PRODUCTION IN TURKEY**

Nazik Meziyet DİLEK, Mustafa KARAKAYA

Selcuk University, Kadir Yallagöz School of Health, Department of Nutrition and Dietetics

[meziyetteemel@hotmail.com](mailto:meziyetteemel@hotmail.com)

**Abstract**

Animal source foods have an important place in human diet, due to the fact that they are rich in terms of digestible protein in high rate. Especially child and young population, that all societies are nourished at the adequate level with animal source foods have a great importance to the physical and mental development. In the studies carried out, for adequate and balanced diet, it was concluded that at least 40 to 50% of daily protein need should be met from animal source foods such as red meat, white meat, milk, and egg. Red meat, one of these foods, is obtained from skeletal muscles of butchery animals and is considerably necessary for a healthy growth and development thanks to high biological value - protein and other dietary members it contains. Being able to meet this need can be enabled society to be able to reach sufficient amount of red meat with minimum cost. Therefore, it is necessary to know the potential of red meat production and the necessity to increase it comes to our face by appropriate breeding policies. In this study, in the light of the last fifteen years, the actual information was attempted to be given regarding the potential of red meat production and animal existence in Turkey.

**Keywords:** Red meat, society, consumption, quality



## NANOTECHNOLOGY IN AGRICULTURE AND FOOD INDUSTRY: AN OVERVIEW OF APPLICATIONS AND CHALLENGES

Rakhshinda Sadiq and Hina Jabeen  
Women University Mardan, Mardan  
[rakhshinda\\_sadiq@yahoo.com](mailto:rakhshinda_sadiq@yahoo.com)

### Abstract

The word nanotechnology is based on the prefix “nano”, the Greek word which means “dwarf”. Furthermore, nanotechnology generally deals with materials having a size range of 0.1 to 100 nanometers resulting in unique physicochemical properties as compared to their macroscopic counterparts. Nanotechnology has a wide range of potential applications in drug delivery and pharmaceuticals. Recently, many scientists and engineers have recognized number of uses of nanotechnology in agriculture and food industries. Use of Nanotechnology based product in agriculture has bright prospects for enhancing the efficacy of agrochemicals and bioavailability of nutrients used through nanoformulations of fertilizers, improving crop yield through bionanotechnology, better control of pests and diseases, , development of a nanoencapsulated pesticide formulation, remediation of contaminated soil and water, improving the shelf-life and quality of fruits vegetables and flowers, etc. The use of nanotechnology also has undoubted benefits for food sector through the use of smart biosensors, packaging materials, and nanonutraceuticals. The benefits of nanotechnology are widespread, as it is a new technology; there are also concerns about toxicity, the impact on human and environmental health. In this review, we intended to cover some of the developments in nanotechnology and their applicability to agriculture and food systems along with potential adverse effects including its legal and regulatory aspects.

**Keywords:** Nanotechnology, agriculture, quality, toxicity

**THREATENED WILD MORELS AND MOUNTAIN LIVELIHOODS IN NORTH WEST PAKISTAN**

Saadia Hanif

Consultant-Farm Dynamics Pakistan (pvt) Ltd.

[saadia.hanif@live.com](mailto:saadia.hanif@live.com)**Abstract**

Rare, short seasoned, wildy grown wild morels are highly valuable edible mushrooms that weigh in gold in worth. Morel exports play a significant role in the socioeconomic condition of mountainous communities of Pakistan. During the months of March to April, picking, drying and selling these forest beauties to exporters is one of major livelihood strategy of these communities specially who are involved in livestock keeping. Wild morels require precise weather conditions to grow and climatic change have resulted in inconsistent precipitation patterns shortening the growing period of morels from two months to one month and in some areas like Swat and Nathiagali, morels have disappeared completely. Deforestation and lack of knowledge related to sustainable practices in morel harvesting has also contributed to the threatening situation for their production and exports. The situation is dire for both local and national economy and ecology. Implementing a carefully designed conserving program involving local communities is highly required. Research programs for creating non timber inventory and preserving forest biodiversity, conducting awareness and training programs in the area for sustainable harvesting techniques for edible forest produce and restrictions over non timber and timber harvests from these forests will help not only saving biodiversity but also sustaining and improving incomes of local households and improving overall economics of the area.

**Keywords:** wild morels, livelihood strategy, sustainable harvest, edible forest produce, forest management

**POLITICAL/IDEOLOGICAL STUDY AND ACADEMIC STUDY:  
SCIENCE,HISTORY AND TEACHING**

Sarmad Hussain Tunio  
Sciences and humanities, NU-FAST Karachi, Pakistan  
[Sarmad.hussain@nu.edu.pk](mailto:Sarmad.hussain@nu.edu.pk)

**Abstract**

There's a lot of difference between the reading of the books with specific political/ideological angle and academic angle. The motives behind the both types of studies are entirely different in our society unfortunately. During the political study, the activists, who already have accepted the told version by the leadership, are mainly concerned with to collect the negative/positive points in order to justify the claims of their leadership regarding the person/event. Political activists are going to perceive the things via certain political lenses, fixed on their eyes by the leadership, which ultimately lead the construction of the biased approach. However, academic study demands objective approach. The academic reader hasn't/shouldn't relatively strong sentiments with the person/event, therefore he/she is only concerned with to understand the things properly. His prime concern is to draw comparatively right conclusion, rather than justifying and then propagating the viewpoint of his leadership. And teaching is another ideological main department but there is not any specific methods that how to teach history, geographphy, science and technology. It is the need of the hour to promote academic traditions, So that social sciences and pedagogical development can be flourished, which ultimately affect the multi dimensional development of our society.

**Keywords:** Ideology,politics,academic study, history science and pedagogy

## THERAPEUTIC EFFECT OF HERBAL PLANTS EXTRACT IN BROILER CHICKEN CHALLENGED WITH COCCIDIA

Subhan Ullah and Umar Sadique

Department of Animal Health, Faculty of Animal Husbandry and veterinary Sciences

The University of Agriculture, Peshawar, Pakistan

[druskhattak@yahoo.com](mailto:druskhattak@yahoo.com)

### ABSTRACT

Coccidiosis is a parasitic disease of poultry birds caused by *Eimeria* species associated with high morbidity, mortality and treatment cost causing a huge economic loss to the poultry industry. The present study was designed to investigate the therapeutic effect of *Azadirachta Indica*, *Curcuma longa* and *Calotropis procera* in broiler challenged with *Eimeria* species. To this end, a total of 360 birds were reared, divided into 6 groups, group A negative control (non-infected and non medicated), group B positive control (infected non-medicated) group C standard group (infected and medicated with Amprolium® @1.25g), group D infected and medicated with *Azadirachta Indica* @ 2g, 4g and 6g, group E infected and medicated with *Curcuma longa* @ 4g, 8g and 12g and group F infected and medicated with *Calotropis procera* @ 4g, 6g and 8g respectively. Group B, C, D, E & F were challenged with 30,000 sporulated oocysts of *Eimeria* species per bird at 21 days and group A was given equal volume of distilled water. Parameters like feed consumption, weight gain, FCR, oocysts in faeces (OPG), mortality and clinical findings were recorded. Current results revealed that the feed intake, weight gain and FCR of non-infected non-medicated group were significantly higher ( $P<0.05$ ) than that of infected non medicated groups. The broiler of group C treated with Amprolium had better significant ( $P<0.05$ ) feed intake and weight gain in comparison to other treated groups. Gross lesions were observed in all medicated groups at low dose level comprising of hemorrhages, congestion and thickening of intestinal wall. Histopathological findings of intestine revealed the sloughing of intestinal villi, leukocytic infiltration and hypertrophy of goblet cell. However no apparent lesions were observed in medicated groups fed with high dose. The result exhibited that *Curcuma longa* @12g/Kg feed had an excellent performance in term of low mortality and oocysts count as compared to other treatment groups.

**Key words:** Broiler, Coccidiosis, Amprolium, *Azadirachta Indica*, *Curcuma longa*, *Calotropis procera*.

## GENOTOXIC AND TOXICOPATHOLOGICAL EFFECT OF AFLATOXIN B<sub>1</sub> IN GRASS CARP (*Ctenopharyngodon idella*)

Hayatullah Khan and Umar Sadique

Department of Animal Health, Faculty of Animal Husbandry and veterinary Sciences

The University of Agriculture, Peshawar, Pakistan

[druskhattak@yahoo.com](mailto:druskhattak@yahoo.com)

### ABSTRACT

Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) is one of the potent mycotoxin produced mainly by *Aspergillus flavus*. Due to the strong mutagenic, teratogenic and carcinogenic effects, AFB<sub>1</sub> is considered as the most common and harmful mycotoxin for animal, human and aquatic life. The present study was designed to evaluate genotoxic and toxicopathological effect of AFB<sub>1</sub> in fresh water fish also known as Grass carp (*Ctenopharyngodon idella*). A total of n=150 Grass carp (42 ± 5 g) were divided into 5 groups each having 3 replicates. Group A was kept as control and group B, C, D and E were exposed to different concentrations of AFB<sub>1</sub> (25, 50, 75 and 100 ppb/kg of diet respectively) for 49 days. *Aspergillus flavus* spores were grown on potato dextrose agar (PDA) slant and then rice was used as a substrate for the production of AFB<sub>1</sub>. The quantity of AFB<sub>1</sub> was 50ppb/g of rice using high performance liquid chromatography (HPLC) technique. The average weight gain and specific growth rate (SGR) of fish was significantly higher in group A (15.35g and 0.63% respectively) as compared to group E (6.55g and 0.28% respectively). The feed conversion ratio (FCR) varied significantly (P<0. 05 between control group and groups containing 75 and 100ppb AFB<sub>1</sub>. AFB<sub>1</sub> did not affect the percent survival rate as no mortality was observed in all treated groups. Furthermore increasing concentrations of AFB<sub>1</sub> significantly lowered the red blood cells (RBC) count, hematocrit, haemoglobin concentration, mean corpuscular volume (MCV), white blood cells (WBC) count and percent lymphocytes as compared to the control group. Similarly a significant increase in the blood biochemical profile such as aspartate amino transferase (AST), alanine amino transferase (ALT), glucose, urea and creatinine was observed in AFB<sub>1</sub> treated groups. The serum protein and albumin level was significantly higher in control group (6.05 & 4.2g/dl) followed by group B (5.8 & 3.9g/dl), group C (5.4 & 3.7g/dl), group D (4.2 & 2.7g/dl) and group E (3.8 & 2.06g/dl). The genotoxicity of AFB<sub>1</sub> was only recorded at 75 and 100 ppb/kg, having micronucleus frequency percentage of 0.85 and 2.15 % respectively. The histopathological study revealed that higher concentrations of AFB<sub>1</sub> can cause pathological changes in liver, kidney, intestine and gills tissue. The present study concluded that AFB<sub>1</sub> causes reduction in fish production performance, effect the haematological and blood biochemical profile, caused tissue damage and induced DNA damage in fish.

**Key Words:** Aflatoxin B<sub>1</sub>, Grass carp, *Ctenopharyngodon idella*, Hematology, Blood Biochemical profile, Histopathology, Micronuclei.

**ANAEMIA – STILL A MAJOR HEALTH PROBLEM**

Binish Sarwar Khan; Syeda Sana Fatima

Institute of Home Sciences, Faculty of Food Nutrition and Home Sciences, University of Agriculture, Faisalabad-Pakistan

Binishasad786@gmail.com; [srizvi665@gmail.com](mailto:srizvi665@gmail.com)

**Abstract**

In Pakistan, approximately 30% boys, 40-50% of primary school and preschool children, 39% of adolescents, 54% girls, 30% of the adult females, 69% children <2 years of age and 47% of the children were reported to affected of Iron deficiency anaemia. The presence of anaemia is basically defined as the lower level of red blood count than the normal range. According to World Health Organization, anaemia is considered as a severe health problem. Like Pakistan, iron deficiency anaemia is also severe health problem in Turkey because the staple food of Turkish people is wheat and 58% of the energy comes from bread with other cereals, which is the major percentage and the average diet from iron is lower than the recommended daily allowance. Women of childbearing age and young children have greater prevalence of iron deficiency anaemia i.e. 40.2% and nearly 2 billion people effects all over the world and nearly one third of the rapidly growing population of approximately 7 billion people. The causes of anaemia were low bioavailability of iron, poor intake of iron, parasitic infections, multiparity and high consumption of those foods that inhibits the absorption of iron, such as tea. For overcoming this severe health problem different six basic implementations have an important role in Pakistan which are: 1) Determining anaemia prevalence 2) Nutrition Training 3) Iron supplementation 4) Controlling parasitic, bacterial or viral diseases 5) Iron fortification in foods 6) Dietary Modifications. To overcome iron deficiency anaemia The Ministry of Health in cooperation with governmental and non-governmental institutions should focus on preventive programs. Nutrition Education is one of the most important prevention program to create awareness among public towards a sound eating habits and healthy lifestyle.

**Keywords:** Anaemia, health, problem, diet, iron

**OCCURRENCE OF *MYCOBACTERIUM BOVIS* IN BOVINE MILK AND CONSUMERS KNOWLEDGE, ATTITUDE AND PRACTICES TOWARD TUBERCULOSIS IN PESHAWAR, PAKISTAN**

Irfan Khattak, Muhammad Hassan Mushtaq, Sultan Ayaz, Sajid Ali, Sadeeq Ur Rahman, Javed Muhammad, Haq Amanullah, Muhammad Luqman Sohail, Furqan Awan, Tahir Usman

College of Veterinary Sciences & Animal Husbandry, Abdul Wali Khan University Mardan, Pakistan

[irfankhattak@awkum.edu.pk](mailto:irfankhattak@awkum.edu.pk)

**Abstract**

This study aims to assess the presence of *Mycobacterium bovis* (*M. bovis*) in retail milk derived from local domestic and commercial dairy farms and the knowledge, attitude and practice about tuberculosis (TB) in the high risk *M. bovis* contaminated milk consumers in Peshawar district, Pakistan. Milk samples were obtained from 92 milk shops and analysed for presence of *M. bovis* using Polymerase Chain Reaction. *Mycobacterium bovis* was detected in 8 milk samples (8.7%). Eight hundred consumers of *M. bovis* contaminated milk were interviewed. Analysis of data indicated that although 97.4% of the participants had heard of TB but only 39.6% knew that cough lasts for more than 3 weeks was one symptom of TB. Only 79.2% had awareness that TB can be prevented and the most frequently stated (48.4%) method of TB prevention was good nutrition. Mean knowledge score for the participants was  $12.1 \pm 2.47$  out of maximum 22. Mean knowledge score varied significantly with ethnicity, level of education and residential status (Urban vs rural). Overall knowledge about symptom, transmission, prevention and treatment of TB was low. Therefore community's health education focused on increasing knowledge of TB must be initiated.

**Keywords:** *Mycobacterium*, milk, dairy farms

**TOXINOTYPING OF *CLOSTRIDIUM PERFRINGENS* INFECTED LAMBS AND KIDS IN DIFFERENT TOPOGRAPHIC AREAS OF KHYBER PAKHTUNKHWA, PAKISTAN**

Mumtaz Ali Khan<sup>1</sup>, Aneela Zameer Durrani<sup>1</sup>, Sher Bahadar<sup>2</sup>, Muhammad Arif Khan<sup>1</sup>, Ali Ahmad Sheikh<sup>1</sup>, Naimat Ullah Khan<sup>3</sup>, Kashif Prince<sup>1</sup>, Naimat Ullah<sup>1</sup> and Ikramullah Khan<sup>2</sup>

<sup>1</sup>University of veterinary and Animal Sciences Lahore

<sup>2</sup>Agriculture University Peshawar

<sup>3</sup>College of Veterinary Sciences and Animal Husbandry Abdul Wali Khan University Mardan

[Dr\\_mumtazkhan@yahoo.com](mailto:Dr_mumtazkhan@yahoo.com)

**Abstract**

In this study project, pathogenic strains of *C. perfringens* were isolated from diseased lambs and kids. The isolates initially identified by colony characters, Gram staining, biochemical tests and CFU/g. The isolates were further analyzed by conventional PCR in order to find out the presence of the alpha, beta, epsilon and iota toxin genes. Pathogen was isolated from 73 of 168 lambs and 86 of 170 kids, having the signs of enterotoxaemia and the animals or their mothers had not been vaccinated previously against *C. perfringens*. Genotyping of 73 strains from diseased lambs indicated (13.10%) prevalence type A, (9.52%) prevalence type B and (20.83%) prevalence of *C. perfringens* type D. Of 86 strains kids (34.71%) prevalence type A, (5.88%) prevalence type B and (10.00%) prevalence of *C. perfringens* type D. *C. perfringens* type C and type E were not detected in both lambs and kids.

Keywords: *C. perfringens*; lambs, kids, toxin genes, PCR



**HEMATO-BIOCHEMICAL ALTERATIONS IN *CLOSTRIDIUM PERFRINGENS* INFECTED GOATS**

Mumtaz Ali Khan<sup>1</sup>, Aneela Zameer Durrani<sup>1</sup>, Sher Bahadar Khan<sup>2</sup>, Muhammad Arif Khan<sup>1</sup>, Ali Ahmad Sheikh<sup>1</sup>, Naimat Ullah Khan<sup>3</sup>, Muhammad Kashif Prince<sup>1</sup>, Naimat Ullah Malik<sup>1</sup> and Azmat Ullah Khan<sup>2</sup>

University of Veterinary and Animal Sciences, Lahore, Pakistan

[dr\\_mumtazkhan@yahoo.com](mailto:dr_mumtazkhan@yahoo.com)

**Abstract**

The study project was designed to determine the affects of *Clostridium perfringens* type D infection on hematological and biochemical parameters in goats. Purposive blood samples were collected from 6 healthy and 12 diseased goats positive for *C. perfringens* infection. Mean erythrocytes count (RBC) and hemoglobin level decreased significantly ( $P < 0.05$ ) in diseased animals while the white blood cells (WBC) increase significantly ( $P < 0.05$ ) in diseased animals compared to the healthy animals. However non-significant differences ( $P > 0.05$ ) were observed in packet cell volume (PCV) and platelets count in healthy and diseased animals. Biochemical analyses indicated significant increases in liver enzymes, total bilirubin, Serum Creatinine, blood Urea and glucose increased significantly ( $P < 0.05$ ) in diseased goats. The results demonstrated that fluctuation in most of the mean hematological values remained within the normal range however the mean liver enzymes, total bilirubin, Serum Creatinine, blood Urea and glucose levels gone beyond the normal levels which demonstrated sever damages to liver and kidneys.

**Keywords:** *Clostridium perfringens*, Goats, RBC, WBC, Platelets, PCV, Bilirubin, Blood Urea, Serum Creatinine.

## QUANTIFICATION, ISOLATION AND PATHOGENOMICS OF PATHOGENIC SHIGA TOXIN ESCHERICHIA COLI O157:H7 ALONG THE PRODUCTION AND SUPPLY CHAIN AROUND HUBEI OF CHINA

Sher Bahadar Khan<sup>1, 2</sup>, Geng Zou<sup>2</sup>, Zia ur Rehman<sup>1, 3</sup>, Farhan Anwar Khan<sup>1, 2</sup>, Atta Muhammad Memon<sup>2, 4</sup>, Ijaz Ahmad<sup>1, 5</sup>, Zahir Shah<sup>1, 2</sup>, Farman Ullah<sup>3</sup>, Shakoor Ahmad<sup>1</sup>, Asad Sultan<sup>6</sup>, Hu<sup>2</sup>, Liu Li<sup>2</sup>, Rui Zhou<sup>2</sup>

<sup>1</sup>Department of Animal Health, The University of Agriculture, Peshawar, Pakistan

<sup>2</sup>Key Lab.of Agriculture Microbiology, Huazhong Agriculture University, Wuhan, China

<sup>3</sup>Key Laboratory of Agricultural Animal Genetics, Breeding and Reproduction, Education Ministry of China, College of Animal Science and Technology, Huazhong Agricultural University, Wuhan 430070, People's Republic of China

<sup>4</sup>Department of Livestock, Govt., of Sindh

<sup>5</sup>National Reference Lab. of Veterinary drug Residues

[sbvetdr@yahoo.com](mailto:sbvetdr@yahoo.com)

### Abstract

Shiga toxin Escherichia coli (STEC) O157:H7 is an important zoonotic food borne pathogens causing gastroenteritis that may lead to life threatening Hemorrhagic Colitis (HC) and Hemorrhagic Uremic Syndrome (HUS). Animal food, particularly poultry, cattle and pig are the major vehicles of diseases caused by food-borne pathogens. A total of 285 samples including 125 tonsil swabs (from 4-6 weeks old clinically healthy animals from different herds of five intensive farms) and 160 samples (20 each liver, meat, intestine and kidney) from different slaughter houses and markets around Hubei province were studied for the prevalence and quantification of STEC O157:H7 along the production and supply chain (PSC). The overall prevalence of E. coli O157:H7 was found 47% along PSC, being higher in supply chain (73.7%) as compared to animal farms (12.8%). Along the supply chain, the highest prevalence was found in slaughter houses (86.25%, 69/80) followed by wet (80%, 32/40) and super markets (42.5%, 17/40). The MPN values ranged from 3-1100MPN/g in overall positive samples, being higher in slaughter houses followed by wet and super markets. Except from intestine and meat samples of slaughter house, the MPN was found higher in summer as compared to winter samples. 8 STEC O157:H7 isolated from meat and liver samples were tested in Balb/C mice for pathogenicity. After development of clinical signs and symptoms, 50-83.3% mortality were produced in the infected mice. Histopathological investigations revealed visible necrosis of intestinal epithelial cells, cell debris shed full of the intestine while in the kidney, renal cortical portion of tubular epithelial cell necrosis were observed. STEC O157:H7 is prevalent along PSC around Hubei of China in different proportions being alarmingly higher in supply chain and markets which is a matter of concern for public health.

**Key words:** Escherichia coli, O157:H7, STEC, MPN PCR, Production, supply chain

## ANTIMICROBIAL RESISTANCE OF BACTERIA ASSOCIATED WITH RAW MILK CONTAMINATED WITH ANTIBIOTICS RESIDUES

Limya .M. Warsma and Ibtisam .M. El Zubeir

Faculty of Animal production dep. of Dairy production, University of Khartoum, Sudan  
[areejwarsmaq3@gmail.com](mailto:areejwarsmaq3@gmail.com)

### Abstract

Milk and other dairy products are produced and processed in Sudan by traditional herdsmen and milk maids. This study was carried out on three hundred raw milk samples (200 cows, 50 camels and 50 goats) that collected from farms, sale points and collection points in Khartoum State during winter and summer season. Milk samples were examined for the presence of antibiotics residues. Also, isolation and identification of bacteria that found in antibiotic contaminated milk and their resistance to some antibiotic residues (ampicillin, cloxacillin, cefotaxime and cephalaxin) were done. Eighty (40%) of cow milk samples were positive to antibiotics residues, while all camel and goat milk samples were negative. Ten (12.5%) and 23 (28.75%) of the positive samples were found during winter and summer, respectively. The positive samples were 47 (58.75%), of which 15 (18.75%) were found during winter and 32 (40%) were during summer. The isolated *Staphylococcus aureus* (2; 0.49%) and *S. auricularis* (4; 0.98%) were sensitive to ampicillin, cephalaxin, cloxacillin and resistant to cefotaxime (75%). *Bacillus cereus* (18; 4.4%) showed resistance to ampicillin, cephalaxin and cloxacillin (100%), while it was sensitive to cefotaxime (93.4%). *Bacillus coagulans* (3; 0.7%) and *B. pumilus* (4; 0.9) were resistant to ampicillin and cloxacillin (100%), however they were sensitive to cephalaxin (33.3% and 50%, respectively). *B. sphaericus* (4; 0.9%) was sensitive to ampicillin (75%), cephalaxin (25%) and cefotaxime (50%) and resistant to cloxacillin (100%). *B. amyloliquefanciens* (3; 0.7%) was resistant to ampicillin and cloxacillin (100%). *Klebsiella* spp. isolates showed sensitivity to ampicillin (33.3%), cephalaxin (100%), cefotaxime (50%) and resistance to cloxacillin (100%). The present study concluded that the quality of milk obtained from cows was lower compared to that of goats and camels. Also, high bacterial loads and antibiotic residues were found in the milk samples collected during summer season compare to winter season.

**Keywords:** Milk, Sudan, bacteria, antibiotics, resistance

**PROCESSING AND UTILIZATION OF TRADITIONAL DAIRY PRODUCTS IN RAWALAKOT AZAD JAMMU AND KASHMIR, PAKISTAN**

Imtiaz Hussain

Department of Food Sciences and Technology, The University of Poonch Rawalakot, AJK Pakistan

[imtiazupr@yahoo.com](mailto:imtiazupr@yahoo.com)**Abstract**

Azad Jammu and Kashmir has diversified traditional dairy product these traditional dairy products can attract the peoples for their unique flavour. The main aim of this study was to assess processing and utilization of traditional dairy products processed in the ruler areas of the Azad Jammu and Kashmir, Pakistan. A single visit multiple subject formal survey was conducted for collection data. A total of 65 households from rural areas of the district Rawalakot, were individually interviewed using a pre designed questionnaire. The major dairy products processed in the targeted area include butter, ghee, boli, kari and Kalri batt. Among these, two of the traditional dairy products, boli, Kalri batt, kari and are unique to this area. Basically, all these traditional dairy products are prepared by simply from the raw milk by fermentation spontaneously at room temperature (17° to 28°C) and then cooked at high temperature (105° to 150°C). The respondents claim that these products are of high nutritional values, hygienically safe and have therapeutic properties. Many aspects of boli, kari, and Kalri batt such as physico-chemical characteristics, microbiological and sensory properties are unknown. Thus, detailed scientific investigation needs to be conducted in order to verify the claimed nutritional and medicinal properties of these products.

**Keywords:** Dairy, processing, utilization, survey

**VALUE ADDITION AND PROCESSED PRODUCTS OF UNDERUTILIZED FRUITS FOR FOOD SECURITY IN AZAD JAMMU AND KASHMIR, PAKISTAN**

Imtiaz Hussain

Department of Food Sciences and Technology, The University of Poonch Rawalakot, AJK Pakistan

[imtiazupr@yahoo.com](mailto:imtiazupr@yahoo.com)

Abstract

Azad Kashmir is endowed with agro biodiversity such as underutilized fruits that are highly nutritious, have health benefits and income generation potential that need to be exploit. These underutilized fruits can be processed in to different products which could contribute to food security and ensure availability of these fruits during off-seasons. The technology should be developed and further disseminated to the community for ensuring food insecurity. These underutilized fruits are important sources of food for rural communities especially at times of food shortage, hunger and other disasters. They provide important compounds for health benefits such as antioxidants, vitamins and minerals. These fruits are considering income generating for rural communities. Different types of food products were processed from these fruits and evaluate for the preference consumer. According to consumer preference test, the juices prepared from apricot and plump have lower value due to its low pH (3.17). The quality of jam, taste and colour were acceptable and preferred. The two products made from fig and pear was all preferred. The microbiological quality of the processed products confirmed their safety characteristics. The prepared products were also evaluated for microbial quality and safety. The findings shows that the products prepared in this experiment were viable, of good quality. Underutilized fruits have great potential to contribute in food and nutritional security.

**Keywords:** Biodiversity, fruit, market, packaging

**HYGIENE AND DAIRY MANAGEMENT PRACTICE DURING MILKING AT RAWALAKOT AZAD JAMMU AND KASHMIR, PAKISTAN**

Imtiaz Hussain

Department of Food Sciences and Technology, The University of Poonch Rawalakot, AJK Pakistan

[imtiazupr@yahoo.com](mailto:imtiazupr@yahoo.com)

**Abstract**

Milk is a complete diet for human it contains numerous nutrients and it makes a significant contribution fulfill the body's needs for minerals, vitamin. If milking is carry out without taking adequate hygienic measures this will cause milk quality deterioration and risk for the consumer health. The main aim of this study was to evaluate some hygienic practices of milking man on dairy farm of Rawalakot. To achieve this objective a study was designed and questionnaires were developed, pre-tested and administered to the selected individuals in the study. About 20 milking personnel and farm attendants related to the selected farms were interviewed. Consequently, hygienic practices employed in the study farms such as house cleaning, udder cleaning, and hand washing practices and milking utensils. The study findings shows that the farmers did not fallow hygiene practice in their routine dairy management and most of the farm hygienic practices and parameters like hygienic condition of the milking environment, sanitation of the milk containers, udder and teats cleaning, use of separate towel for each cow and the personal hygiene of the milkers were not fully performed by most of the farm owners. The farmers used a single cloth in washing the udder for several animals and did not thoroughly wash their hands during milking. Low grade plastic cans and bottles which could cause quality deterioration were used in milking and delivering milk to the collection points and consumer.

**Keywords:** Milk, management, hygiene

## NUTRACEUTICAL APPLICATIONS OF ZINC, MANGANESE AND COPPER IN THE METABOLISM OF THE RUMEN AND IMMUNE FUNCTION

Habib-Ur-Rehman1\*, Fazal ur Reman1, Ali akbar1, Munir Ahmad1, Niamatullah Kakar2, Zafar Ahmad2, Asmatullah Kakar3, Muhammad Arif Kakar4, Qudratullah5, Mir Ahmad6, Muhammad Ayub7 and Siraj Ahmad7.

Department of Microbiology, University of Balochistan (UoB), Quetta, Pakistan

[hur\\_2085qta@yahoo.com](mailto:hur_2085qta@yahoo.com)

### Abstract

Balance among the essential nutrients, protein, carbohydrates, fats, minerals and vitamins is a key component in striving towards optimum animal production. The role of trace minerals in animal production is an area of strong interest, as the trace elements recognized the most important functional components of numerous metabolic events. Trace minerals are of basic needs by the body in trace amounts, ranging from 0.10 to 50.0 mg/kg of dry matter in the feed of bovine animals. These trace elements are essential for all biochemical processes of the body that supports the growth and the appropriate maintenance. For example, copper is necessary for the function of the superoxide dismutase and in the elimination of toxic by-products of metabolic pathways. The elimination of these toxic by-products allows metabolism to perform effectively, uninhibited by damaging oxygen free radicals. Manganese, an important trace mineral, essential for development, metabolism, in the antioxidant system and has a slight effect on stimulating the activity of the Urease. Zinc, an important trace mineral for the enzymatic function, help in the regulation of the production of nucleic acid, the metabolism of carbohydrates and protein synthesis, thus providing a stable framework for the development. The immune system is part of the defense of the host against the destructive forces from the outside of the body, such as bacteria, fungi, parasites and viruses, or of the Interior, such as the malignant cells or those who produce auto-antibodies. Trace minerals have an important role for normal immune function and disease resistance including zinc, copper and manganese. An insufficiency in at least one of these components can compromise immunocompetence of an animal. The immune framework is made of two branches: the innate or non-specific of the immune framework, and the adaptive or specific framework of immunity. In this review paper, an effort has been established to scrutinize the effects of minerals supplement in the metabolism of rumen and their individual or joined effects on immune function in different species of animals.

**Keywords:** Minerals, copper, manganese, zinc, metabolism, ruminants

## REUSE OF WASTE WATER IN AGRICULTURE RECOVERED FROM WASTEWATER TREATMENT SYSTEMS

Khush Bakht Andleeb & Imran Hashmi

Institute of Environmental Sciences and Engineering: School of Civil and Environmental Engineering, Islamabad, Pakistan. National University of Sciences and Technology, Islamabad, Pakistan

[kbandleeb@gmail.com](mailto:kbandleeb@gmail.com)

In adequate access to sanitation and clean water has become one of the most pervasive problems affecting people throughout the developing world. Conventional wastewater treatment technologies have proved ineffective in solving complex water related issues resulting from rapid industrialization and urbanization throughout the developing world. Membrane bioreactors (MBR) and Phytoremediation system under study showed to be very robust Eco technologies for domestic wastewater treatment and reuse. The present study was conducted at NUST H-12 campus Islamabad to compare the water quality of MBR and Phytoremediation system in terms of Physico-chemical and Biological parameters and microbial species present in the system. Phytoremediation system under study was planted with *Typha latifolia*, *Pistia stratiotes*, *Centella asiatica*. An excellent overall treatment performance was exhibited on the conventional water quality parameters (up to 90.8% for TSS, 67.7% for COD, 95% for turbidity and 90% for total coliform) by Phytoremediation system. Treatment performance of MBR system was up to 99% for TSS, 83% for COD, 91.5% for turbidity and 99% for total coliform and final effluents proved to compile with the EPA regulations. The high removal rates were achieved at higher temperature as well as other meteorological parameters (wind speed, air pressure, relative humidity, global horizontal irradiance) showed a significant positive and negative correlation with the removal efficiencies. Predominant species isolated and identified from wastewater of Phytoremediation system belongs to the phyla Proteobacterium (*Enterobacter cloacae*, *Enterobacter kobei*, *Enterobacter hormaechei*, *Enterobacter asburiae*, *Enterobacter aerogenes*, gamma proteobacterium, *Franconibacter pulveris*, *Citrobacter freundii*, *Shigella dysenteriae*, *Escherichia albertii* and *Escherichia coli*). While, predominant bacterial species isolated and identified from activated sludge of MBR system were *Salmonella enterica*, *Pantoea dispersa*, *Shigella dysenteriae*, *Enterobacter hormaechei* and *Salmonella waycross* and they too belong to the phyla proteobacteria. High efficiency of these treatment systems will cater the water need of agricultural land and divert the path of fresh water to drinking purposes.

**Keywords:** wastewater, agriculture, treatment systems



**CLINICAL & RESEARCH LABORATORIES IN PESHAWAR CITY OF KP PROVINCE: ANALYSIS OF BIOSECURITY MEASURES PREDICT INSECURITY**

Hoor Shumail and Tayyab Rehman

Department of molecular biology & genetics, Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, Pakistan

[hoorshumail.bioinformatics@kmu.edu.pk](mailto:hoorshumail.bioinformatics@kmu.edu.pk)

**Abstract**

Biosecurity is an emerging concept not well understood by the local community. It is the protection, control and accountability of biological agents and toxins within laboratories, in order to prevent their loss, theft, misuse, unauthorized access or intentional unauthorized release. All types of wet laboratories are responsible for biosafety and biosecurity that protects their workers, animal population, and the environment from exposure or spread of pathogens they come across with. The purpose of this preliminary study was to collect data in order to assess and evaluate beside biosafety practices the measures being taken for biosecurity in clinical diagnostic and research lab. In this study hundreds of clinical labs, Research labs and Academic purpose labs for routine practical work both in public and private setups were focused. The visits to the sites were pre scheduled and in some cases were informal and unscheduled with the hope to get a clear picture of the practices/measures. A detailed observational survey and unwritten questions were used to elicit information. The focus was on laboratory licensing system, infrastructure, administration, quality assurance, collection, handling, processing, storage and transportation of clinical samples, waste disposal, Lab records, access to biohazardous materials and security checks on pathogens and on researchers and workers. Based on Standard BRM Guidelines, maximum numbers of the labs were below minimum required standard and were categorized as BSL-0. Variety of BSL-2 and BSL-3 category bacterial and viral pathogens were easily assessable and, therefore, were isolated and identified in our lab. We conclude that lab management is lacking at all levels. It is recommended that biosecurity concerns be conveyed to all stakeholders for designing and implementing action plan.

## ECOLOGICAL BASED FEED INVENTIONS FOR LAMBS/KIDS FATTENING MANEUVER TO EARN GREATER PROFIT REVENUES DURING EID-UL-AZHA MARKETING IN PAKISTAN

Abdul Razzaq<sup>1</sup>, Muhammad Islam<sup>1</sup>, B. Ann Rischkowsky<sup>1</sup>, Ataullah Khan<sup>2</sup>, Gul Muhammad<sup>2</sup>, M.N.M. Ibrahim<sup>3</sup>, Sarfraz Ahmad<sup>4</sup> and Farman Ullah<sup>5</sup>

<sup>1</sup>International Center for Agriculture Research in the Dry Areas (ICARDA), Jordan

<sup>2</sup>Arid Zone Research Institute, PARC, Umerkot

<sup>3</sup>International Livestock Research Institute (ILRI), Nairobi, Kenya

<sup>4</sup>Pakistan Agricultural Research Council

<sup>5</sup>Faculty of Veterinary and Animal Sciences, Lasbela University of Agriculture, Water and Marine Sciences, Uthal, Balochistan, Pakistan

[abdulrazzaqzrc@gmail.com](mailto:abdulrazzaqzrc@gmail.com)

The main aim of this study was to determine the impact of supplemental feeding on lamb/kids fattening based on feed availability in three ecologies during Eid-UI-Azha marketing of Pakistan. This is the special occasion where farmers can get more than double profit than rest of the year marketing. For this purpose, private farmers (3-5) from each of three research sites namely; Dhulli village at district Chakwal in Punjab, Loralai in Balochistan and Kunri village at Umerkot in Sindh were selected for the fattening trials. Three feed formulations i.e., T1, T2 and T3 were self-prepared as per availability of ingredients for Chakwal site experiment. However two commercial rations T4 and T5 used for Loralai site and other two commercial rations T6 and T7 for Umerkot site as per availability. The estimated Crude Protein (CP), Total Digestible Nutrients (TDN) and Energy (MJ/kg) in these formulations were ranged between 15-17.85%, 70-91% and 2800-3200 respectively. The overall results revealed that lambs/kids attained higher live-weight gain (2.4 to 8.5 kg: 96-188 g/head/day) with supplemental feeding compared with farmers practices (0.7 to 2.8 kg: 23-62 g/head/day). The overall economic analysis showed that lambs/kids fed with supplemental rations leads to higher profit margins which ranged from Rs. 5190/- to Rs. 14730/- per head compared with control groups (Rs. 5100/- to Rs. 12900/-). The results revealed that with supplemental feeding Rs. 90/- to 1830/- can be earned higher compared to farmer's practices. Therefore, it is recommended that area specified feed supplement options may be utilized for higher profit earning specially during the Eid-UI-Azha marketing in Pakistan.

**Keywords:** Ecological based feeding, small ruminants, fattening, Eid-UI-Azha marketing

## IDENTIFICATION OF NON-GENETIC FACTORS AFFECTING WEIGHTS AT BIRTH, PRE-POST WEANING, AND GREASY FLEECE PERFORMANCE OF KAJLI SHEEP IN PAKISTAN

Farman Ullah<sup>1</sup>, Khalid Javed<sup>2</sup>, Mohammad Salim<sup>3</sup>, Momen Khan<sup>4</sup>, Sajid Ali<sup>5</sup>, Abdul Razzaq<sup>6\*</sup>, Illah Bakhsh Marghazani<sup>1</sup>, Muhammad Altaf Hussain<sup>1</sup>, Ziaurrahman<sup>7</sup>, Sibghatullah<sup>8</sup>, Tariq Zeb<sup>9</sup>, Dinesh Bhattiarai<sup>7</sup>, Talpur Hira Sajjad<sup>7</sup>

<sup>1</sup>Faculty of Veterinary and Animal Sciences, Lasbela University of Agriculture, Water and Marine Sciences, Uthal, Balochistan, Pakistan; <sup>2</sup>Faculty of Animal Production and Technology, University of Veterinary and Animal Sciences, Lahore, Punjab, Pakistan; <sup>3</sup>Department of Forestry and Wildlife Management, University of Haripur, Khyber Pakhtunkhwa, Pakistan; <sup>4</sup>Livestock and Dairy Development Department, Khyber Pakhtunkhwa, Pakistan; <sup>5</sup>Agricultural university Peshawar Amir Muhammad Khan Campus Mardan; <sup>6</sup>Animal Sciences Division, Pakistan Agricultural Research Council, Ataturk Avenue, Islamabad; <sup>7</sup>Key Laboratory of Agricultural Animal Genetics, Breeding and Reproduction, Education Ministry of China, College of Animal Science and Technology, Huazhong Agricultural University, Wuhan 430070, People's Republic of China; <sup>8</sup>Foot and mouth disease research center, Veterinary research institute Peshawar; <sup>9</sup>Veterinary Researches and Disease Investigation Center, Swat, Pakistan

[abdulrazzaqzrc@gmail.com](mailto:abdulrazzaqzrc@gmail.com)

### Abstract

The present investigations were to analyze the influence of environmental factors on various growth performance traits of Kajli sheep in Pakistan. The present study utilizes two large contemporary datasets in sheep to explore factors that influence performance traits. The study was conducted to estimate the non-genetic components of variation influencing various traits of economic importance in Kajli sheep. For this purpose reproduction performance recorded of 13715 Kajli sheep lambing collected during 1994 to 2010 at Livestock Experimental Stations Khushab and Khizarabad, Punjab. The data on Birth Weight (BW), 120 days at weaning Weight (WW), Pre-weaning Average Daily Gain (PRADG), Yearling Weight (YW) and Greasy Fleece Weight (GFW) were evaluated. Location of flock, year of birth (YOB), Birth Season (BS), Birth Types (BT), and sex were the fixed effects in the model. Results showed that, the overall average values for birth weight (kg), weaning weight (kg), yearling weight (kg), pre weaning weight (gm), and greasy fleece weight (kg) were  $4.13 \pm 0.01$ ,  $18.70 \pm 0.08$ ,  $37.52 \pm 0.06$ ,  $142.34 \pm 0.83$  and  $1.32 \pm 0.00$ , respectively. Year of birth, type of birth, sex and flock influenced ( $P < 0.001$ ) birth weight and greasy fleece weight whilst season of birth showed no significant differences ( $P > 0.05$ ). In weaning weight and pre weaning average daily gain of Kajli sheep, year of birth, type of birth and flock showed influence ( $P < 0.05$ ) except sex ( $P > 0.05$ ). In yearling weight, all parameters showed effect ( $P < 0.05$ ) except type of birth and flock ( $P > 0.05$ ). The difference might be attributed to regular supplementation of ration to flock. Male lambs were heavier than female lambs, and single born lambs were also significantly heavier than twins ( $P < 0.05$ ) at birth. Results reflected that Kajli breed can be improved through selection and better management.

**Keywords:** Kajli Sheep, Environmental factors, performance traits, variation, Pakistan

**(Addition of more abstracts in process)**

## Authors Index

- Acar, Gamze ; EFFECTS OF OLIVE LEAVES ON SHELF LIFE ; SU Konya Turkey; 091  
Agossou, Dehouegnon Jerry; THERMAL AND NUTRITIONAL; Çukurova Univ; Konya, Turkey; 040  
Ahmad, Ishtiaque; ANTIMICROBIAL POTENTIAL OF; UVAS Lahore, Pakistan; 016  
Ahmad, Maqbool; COMPARATIVE EFFICACY OF SYNCH; UAF, Faisalabad, Pakistan; 018  
Ali, M; ESTIMATION OF COMPOSITION, ADULTERATION; UVAS, Lahore, Pakistan; 005  
Ali, Muna Jalal ; ANTIMICROBIAL PATTERN COMBINED ACTION ;UMP Malaysia ; 084  
Alp, Halime; SOME TECHNOLOGICAL PROPERTIES OF MEAT PROTEIN; SU Konya Turkey; 081  
Andleeb, Khush Bakht; REUSE OF WASTEWATER IN AGRICULTURE; NUST Pakistan; 068  
Ashraf, M; ANTHELMINTIC AND ANTIPROTOZOAL; UVAS, Lahore, Pakistan; 006  
Babaoğlu, A S; CHARACTERISTICS OF MEAT EMULSION; SU, Konya, Turkey; 010  
Bakhtawar, Maqbool; DIFFERENTIAL GROWTH OF SALMONELLA, UAF, Faisalabad, Pakistan; 013  
Batool, Maryam ; COLOSTRUM WHEY: A POTENTIAL; UVAS, Lahore, Pakistan; 012  
Chand, N. ; EFFECT OF FENUGREEK SEED ; AUP Pakistan; 047  
DİLEK, Nazik Meziyet ; CONJUGATED LINOLEIC ACID IN ANIMAL; SU Konya; 050  
DİLEK, Nazik Meziyet ; IDENTIFICATION OF PROCESSED MEAT PRODUCTS ; SU Konya; 051  
DİLEK, Nazik Meziyet ; POTENTIAL OF RED MEAT PRODUCTION ; SU Konya 052  
DİLEK, Nazik Meziyet; DETERMINATION OF RED AND WHITE MEAT ;SU Konya; 048  
DİLEK, Nazik Meziyet; IDENTIFICATION OF THE PREFERENCES; SU Konya; 049  
EFFECT OF RIPENING CONDITIONS ON PHYSICO-CHEMICAL ; UVAS, Lahore, Pakistan 024  
Farmanullah ; POINT PREVALENCE OF GASTROINTESTINAL; LUA Balochistan Pakistan ; 098  
Farmanullah ; UPSHOT OF INHERITED FACTORS; HAU China; 104  
Gul, Shafia Tehseen ; DIAGNOSIS OF LATENT BACTERIAL; UAF Faisalabad Pakistan; 089  
Hanif , Nafeesa Qudsia ; HOMO FERMENTATIVE BACTERIAL; AgriPak Labs, Pakistan; 074  
Hanif, Saadia; THREATENED WILD MORELS AND; Farm Dynamics Pakistan (pvt) Ltd ; 054  
Hashmi, Imran; INVESTIGATING DISINFECTION BY-PRODUCTS ; NUST Islamabad Pakistan; 079  
Hayat, Shams-ul; INCIDENCE RATE OF BLOOD PROTOZOANS; VR&DIC, Kohat ;103  
Hoor Shumail and Tayyab Rehman; CLINICAL & RESEARCH; KMU; Peshawar Pakistan; 069  
Hosseini, Seyed Mahdi; DETERMINATION OF BREEDING; AREEO, Qom, Iran; 085  
Hussain, Imtiaz ; HYGIENE AND DAIRY MANAGEMENT PRACTICE ; UoP Azad Kashmir; 066  
Hussain, Imtiaz ; VALUE ADDITION AND PROCESSED PRODUCTS ; UoP Azad Kashmir; 065  
Hussain, Imtiaz; PROCESSING AND UTILIZATION OF TRADITIONAL; UoP Azad Kashmir ; 064  
Hussain, Jazib; COMPARATIVE EVALUATION OF NIGELLA SATIVA; UAF, Pakistan 043  
Hussain, Jazib; COMPARATIVE EVALUATION OF NIGELLA; UAF, Faisalabad, Pakistan ; 036

Ikram, M Asim; QUALITY ASSESMENT OF PASTEURIZED ; UVAS, Lahore, Pakistan ; 003  
 Imran,M ; IMPACT OF GRADED LEVELS OF INOCULANT ; QAU Islamabad Pakistan; 078  
 Inam, M; EFFECT OF SACCHAROMYCES CEREVISIAE; SBBU Shringal Pakistan; 100  
 Iqbal, Aamir; EFFECT OF MENTHA PIPERITA (PEPPERMINT; UVAS, Lahore-Pakistan ; 039  
 Iqbal, Zahid; DISPOSITION KINETICS OF CIPROFLOXACIN; Isra University; 008  
 Ismail, Amir ; A REVIEW ON AFLATOXIN M1; BZU, Multan, Pakistan ; 011  
 Jabeen, Hina; PHYTOREMEDIATION: AN INSPIRING; WU Mardan, Pakistan; 042  
 Jamal, Muhammad; WATER PRODUCTIVITY UNDER; UAP, Peshawar, Pakistan ; 019  
 Jan, Amanullah ; ANIMAL BASED ORGANIC CARBON SOURCES; UAP Peshawar Pakistan;  
 090  
 Jan, Awsi; EFFECT OF GAMMA-IRRADIATION; SKUAST, Jammu and Kashmir; 033  
 Jawad, Muhammad Aslam, MONOGLYCERIDES AN ALTERNATIVE TO; UAF Pakistan; 046  
 Kakar, M Azam; CRIMEAN CONGO HEMORRHAGIC FEVER; UAP, Peshawar, Pakistan ;  
 002  
 Kakar, M.A. ; CRIMEAN CONGO HEMORRHAGIC FEVER (CCHF) ;UAP Peshawar  
 Pakistan; 096  
 Kausar, Razia ; ESTIMATION OF MILK PROTEASE ACTIVITY ; UAF, Faisalabad, Pakistan ;  
 037  
 Kausar, Razia ; ESTIMATION OF MILK PROTEASE ACTIVITY ;UAF Faisalabad; 087  
 Khan , Mumtaz Ali ; HEMATO-BIOCHEMICAL ALTERATIONS ; UVAS Lahore Pakistan;  
 061  
 Khan , Mumtaz Ali; TOXINOTYPING OF CLOSTRIDIUM PERFRINGENS; UVAS Pakistan;  
 060  
 Khan, Aman Ullah ; SEROPREVALENCE AND POTENTIAL RISK; UVAS ,Lahore, Pakistan ;  
 001  
 Khan, Amjad; EPIDEMIOLOGY OF ZONOTIC BRUCELLOSIS ; UVAS Lahore Pakistan;  
 102  
 Khan, Mir Usman; ENHANCEMENT OF OXIDATIVE STABILITY; UVAS Lahore, Pakistan ;  
 026  
 Khan, Mir Usman; PHYSICO-CHEMICAL AND NUTRITIONAL ; UVAS Lahore, Pakistan;  
 025  
 Khan, Sher Bahadar; QUANTIFICATION, ISOLATION AND PATHOGENOMICS; UAP  
 Pakistan ;062  
 Khattak, Irfan; OCCURRENCE OF MYCOBACTERIUM BOVIS; AWKUM Mardan Pakistan;  
 075  
 Khattak, Irfan; OCCURRENCE OF MYCOBACTERIUM BOVIS; AWKUM Pakistan; 059  
 M Imran ; OF GRADED LEVELS OF INOCULANT ; QAU Islamabad Pakistan; 099  
 Mahmood, Aisha; OMEGA-6/OMEGA-3 RATIO IN MARGARINE; UAF, Faisalabad, Pakistan  
 ; 032  
 Malik, Saleem; APPLICATIONS OF PULSE ELECTRIC; SKUAS, Jammu and Kashmir; 038  
 Masood, Zubia; AN INVESTIGATION ON CHEMICAL; JUW, Karachi, Pakistan; 029  
 Masoodi, Lubna; EFFECT OF ANTIOXIDANT, PACKAGING; SKUAST, Srinagar, J&K; 044  
 Muzaffar, Humaira ; HIGH PROTEIN DIET INDUCES; UAF, Faisalabad, Pakistan ; 015  
 Naveed, Ahsan ; COMPARATIVE EVALUATION OF; UAF, Faisalabad, Pakistan ;031  
 Nazeer, Hamza; COMPARISON OF LINCOMYCIN AND; UAF, Faisalabad, Pakistan; 041  
 NidaUllah, Hafiz; RISK CHARACTERIZATION OF ANTIBIOTIC; Universiti Sains Malaysia;  
 035  
 Nizam, Maria; AN ANALYSIS OF CORRELATION BETWEEN POOR ; GU Peshawar  
 Pakistan; 045

Okaka, Wilson ; PUBLIC AWARENESS COMMUNICATION; Kyambogo University, Uganda ; 028

Okaka, Wilson; STRENGTHENING COMMUNITY RADIO FOR; Makerere University, Uganda ; 027

Omer, Muhammad Ovais; IN-VITRO CYTOTOXIC AND ANTIVIRAL; UVAS Lahore Pakistan; 095

Qamar, M Fiaz; CHALLENGES AND FUTURE SCOPE; UVAS, Lahore, Pakistan ; 004

Qureshi, Irfanul Haq; DETERMINANTS OF LABOR FORCE; IMS , Peshawar Pakistan; 073

Qureshi, Samia Subhan; POSITIONING AND NAVIGATION; UET Peshawar Pakistan; 083

Qureshi, Sania Subhan; DAIRY SCIENCE PARK–EMPLOYMENT; UAP, Peshawar, Pakistan ; 023

Raja, Rameez; EFFECT OF BREED ON CARCASS TRAITS; SAU, Tandojam, Pakistan 022

Razzaq, Abdul; ECOLOGICAL BASED FEED INVENTIONS FOR LAMBS; ICARDA, Jordan; 070

Razzaq, Abdul; IDENTIFICATION OF NON-GENETIC FACTORS; ICARDA, Jordan; 071

Rehman Habib-Ur ; THE SIGNIFICANCE OF SELENIUM ; UoB Quetta Pakistan; 101

Rehman, Habib Ur; NUTRACEUTICAL APPLICATIONS OF ZINC ; UoB Quetta, Pakistan ; 067

Rehman, Tayyab ur; FROM DAIRY SCIENCE PARK TO; KMU Peshawar Pakistan ; 082

RIZWAN, H.M. ; SYNTHETIC POLYMORPHIC PEPTIDES ; UAF, Faisalabad, Pakistan ; 020

Rizwan, M Hafiz; FASCIOSIS: A MAJOR THREAT TO FOOD; UAF, Faisalabad, Pakistan ; 021

Saddique , Umer; GENOTOXIC AND TOXICOPATHOLOGICAL EFFECT; UAP Pakistan; 057

Saddique, Umer; GENOTOXIC AND TOXICOPATHOLOGICAL; UAP Peshawar Pakistan ; 094

Sadiq, Rakhshinda ; NANOTECHNOLOGY IN AGRICULTURE ; WU, Mardan ; 053

Sadique, Umar ; THERAPEUTIC EFFECT OF HERBAL PLANTS; AUP; 056

Salih, Barka Mohammed; PREVALENCE OF ESCHERICHIA; Route de la Rocade, Algeria; 034

Sarwar, Binish Khan; ANAEMIA – STILL A MAJOR HEALTH PROBLEM; UAF Pakistan; 058

Shah, Zahir ; BIOFERTILIZERS FOR SUSTAINABLE; AUP Peshawar Pakistan ; 086

Shamim, Asim ; COMPARATIVE RESILIENCE OF TEDDY; UOP, Azad J & Kashmir, Pakistan; 014

Sohail, Sajid Muahammad; EPIDEMIOLOGY AND CONTROL; UAF, Faisalabad, Pakistan ; 030

Suhail, Syed M ; EFFECT OF YEAST CULTURE (SACCHROMYCES; AUP Peshawar Pakistan; 076

Suhail, Syed M ; STUDY ON GROWTH PARAMETERS; AUP Peshawar Pakistan; 077

Sultan, Asad ; TOTAL TRACT NUTRIENT RETENTION ; UAP Peshawar Pakistan ; 097

Tufail, Muhammad Shoaib; IMPACTS OF USING IMPROVED SEED OF; CSU Australia; 088

Tunio, Sarmad Hussain ; POLITICAL/IDEOLOGICAL STUDY; NU-FAST Karachi; 055

Uddin, Rafi; EFFECT OF REPLACEMENT OF MAIZE; L&DD Balochistan Pakistan; 080

Ünal, Kübra; THE EFFECTS OF KEFIR IN TENDERIZING; Selçuk University, Konya, Turkey; 017

Usman, Tahir ; TWO NOVEL SNPS IN STAT3 GENE ARE; AWKUM, Mardan, Pakistan; 092

Waqas, M Khurram; OCCURANCE AND CAUSES; UVAS, Lahore, Pakistan; 093

Warsma, Limya .M.; ANTIMICROBIAL RESISTANCE OF BACTERIA; UoK Sudan; 63

Younas, M; TOXICOLOGICAL EFFECTS OF AFLATOXIN B1; UVAS, Lahore, Pakistan; 007



Zia ur Rehman; EPIDEMIOLOGICAL RISK FACTORS ; HAU Republic of China; 072  
Zubair, M; EFFECTS OF DIETARY VITAMIN C; University of Ponch, Azad Kashmir ; 009



# Participating organizations

## International

1. [DSP-UN Partnership – UNDESA New York](#)
2. [Chinese Academy of Agricultural Sciences, \(CASS\) China](#)
3. [Charles Sturt University Australia](#)
4. [International Livestock Research Institute Kenya](#)
5. [Engormix.com Argentina](#)
6. [The International Society of Zoological Sciences China](#)
7. [Selcuk University Konya, Turkey](#)
8. [Sandia National Laboratories USA](#)
9. [Smith and Franklins Publishers UK](#)
10. [Teknokent Konya Turkey](#)
11. [Highway Conference Discovery System China](#)
12. [SETBIR Turkey](#)
13. [ASUD Turkey](#)
14. [HICDS China](#)
15. [Amanah Halal Research Centre Germany](#)

## National – Corporate

1. DSP Slaughtering Point, Bar Tehkal, Peshawar
2. Jan Arcade Meat Shop, Peshawar
3. Meat Park, Peshawar
4. Dairy Life, University Feed Mill
5. Mr Kamal Khan, silage industry
6. Reon Energy Solutions, Lahore
7. Ghazi Brothers, Lahore
8. DSP Innovations
9. Farmer's Sausage Company, Peshawar
10. Royal Dairies, Peshawar
11. CATTLEKIT, Lahore
12. Livestock Exporters Assoc Pak
13. Riaz Park, Bannu
14. Romer Labs, Pakistan, Rawalpindi
15. Asif and Co, Peshawar
16. Calves and Calves Pvt Ltd
17. ICI Pakistan Ltd
18. M Dairy Solutions Pakistan
19. Solve Agric Pakistan Ltd
20. Globe group of companies
21. Peshawar PCR Lab (PPL), Emergency Gate LRH, Peshawar

## National – Development

1. SDGs Task Force, Khyber Pakhtunkhwa Provincial Assembly
2. [Sustainable Development Unit, Govt of KP](#)
3. Small and Medium Enterprises Development Authority Peshawar
4. [Khyber Medical University, Peshawar](#)
5. [University of Agriculture Peshawar](#)
6. Khyber Institute of Veterinary Sciences, Peshawar
7. SBBU, Sheringal
8. Veterinary Research Institute, Peshawar
9. CEBG, Military Farms Group Okara
10. [Women University Mardan](#)
11. [Halal Research Council Lahore](#)
12. [Banking – the exclusive bimonthly magazine on Islamic Banking and Finance](#)